

CENTRAL LIMIT THEOREMS FOR ITERATED RANDOM LIPSCHITZ MAPPINGS

HENNION Hubert¹ and HERVE Loïc²

I.R.M.A.R., Laboratoire de Probabilités, Université de Rennes 1.

Abstract. Let M be a non-compact metric space in which every closed ball is compact, and let G be a semi-group of Lipschitz mappings of M . Denote by $(Y_n)_{n \geq 1}$ a sequence of independent G -valued, identically distributed random variables (r.v.), and by Z a M -valued r.v. which is independent of the r.v. Y_n , $n \geq 1$. We consider the Markov chain $(Z_n)_{n \geq 0}$ with state space M which is defined recursively by $Z_0 = Z$ and $Z_{n+1} = Y_{n+1}Z_n$ for $n \geq 0$. Let ξ be a real valued function on $G \times M$. The aim of this paper is to prove central limit theorems for the sequence of r.v. $(\xi(Y_n, Z_{n-1}))_{n \geq 1}$. The main hypothesis is a condition of contraction in the mean for the action on M of the mappings Y_n ; we use a spectral method based on a quasi-compactness property of the transition probability of the chain mentioned above, and on a special perturbation theorem.

SHORT TITLE : LIMIT THEOREMS FOR ITERATED MAPS

AMS subject classification : 60J05-60F05

Keywords : Markov chains, iterated function systems, central limit theorems, spectral method.

¹Université de Rennes 1, Campus de Beaulieu, 35042 Rennes Cedex. Hubert.Hennion@univ-rennes1.fr

²Institut National de Sciences Appliquées de Rennes, 20, Avenue des Buttes de Couësmes CS 14 315, 35043 Rennes Cedex. Loic.Herve@insa-rennes.fr

1 Introduction

Let M be a non-compact metric space in which every closed ball is compact, endowed with its Borel σ -field \mathcal{M} . We denote by G a semi-group of Lipschitz mappings of M and by \mathcal{G} a σ -field on G . We assume that the action of G on M is measurable, i.e. the map j defined by $j(g, y) = gy$ is measurable from $(G \times M, \mathcal{G} \otimes \mathcal{M})$ to (M, \mathcal{M}) .

Let π be a probability distribution on G , and let $(Y_n)_{n \geq 1}$ be a sequence of independent G -valued random variables (r.v.) identically distributed according to π , defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The iterated random mappings R_n , $n \geq 0$, are defined by

$$R_0 = Id_M, \quad R_n = Y_n \cdots Y_1, \quad n \geq 1.$$

Let Z be a M -valued r.v. which is independent of the r.v. Y_n , $n \geq 1$. The sequence $(Z_n)_{n \geq 0}$ defined by

$$Z_n = R_n Z, \quad n \geq 0$$

is a Markov chain on M which is defined recursively by

$$Z_0 = Z, \quad Z_{n+1} = j(Y_{n+1}, Z_n) = Y_{n+1} Z_n, \quad n \geq 0.$$

Observe that we get here the general Lipschitz iterative model on M which has been considered by many authors, see Duflo (1997), Diaconis and Freedman (1999) to get an overview of the subject. Consider particularly the case where M is the linear space \mathbb{R}^q . The generalized linear autoregressive model is obtained when G is the semi-group of affine mappings of M . Replace in the preceding the linear part of the action by that of a fixed Lipschitz mapping f of M . An element g of the semi-group G is now defined by a vector $b_g \in M$, and it acts on M according to the formula $gx = f(x) + b_g$. In this context the probability distribution π on G is simply defined by a distribution on M , thus we get the Lipschitz functional autoregressive model.

Now let ξ be a real valued measurable function on $G \times M$. The aim of this paper is to establish a central limit theorem with a rate of convergence and a local central limit theorem for the sequence of r.v.

$$(\xi(Y_n, Z_{n-1}))_{n \geq 1}.$$

The interest of considering a function ξ of the couple $(g, x) \in G \times M$ rather than a function only depending on x appears for example in the study of random matrices products.

From the stochastic view point, the context may be described as the study of the sequence of r.v. obtained by composing the function ξ and the Markov chain $(X_n)_{n \geq 0}$ with state space $G \times M$ defined by

$$X_0 = (Id_M, Z), \quad X_n = (Y_n, Z_{n-1}), \quad n \geq 1.$$

The main hypothesis will be a condition of contraction in the mean of the action on M of the elements of G under the probability distribution π . This property enables us to make use

of a refinement of the spectral method. Recall that the spectral method has been initiated by Nagaev (1957), and then used and improved by many authors. It is fully described in Hennion and Hervé (2001) where references are given. The spectral method is based on a quasi-compactness property of the transition probability Q of the chain $(X_n)_{n \geq 0}$, and on a perturbation theorem ensuring that, for small $|t|$, the Fourier kernels $Q(t)$ associated with Q and ξ have spectral properties similar to those of Q . In the present setting, the use of the standard perturbation theory for operators leads to assume moments of exponential type (cf. Milhaud and Raugi (1989), Hennion and Hervé (2001) Chap. X, Sect 3). The main feature of this paper is the use of a perturbation theorem of Keller and Liverani (1999) which is adapted to operators verifying a Doeblin-Fortet inequality. By means of this theorem we get the desired limit theorems under moments of polynomial types.

Notice that there are several methods to cope with central limit theorems for a function of a Markov chain, mostly known are regeneration and splitting, use of central limit theorems for martingale increments and Lindeberg techniques. As will be discussed later, when applied to the present context, some of these methods can give a central limit theorem under hypotheses which are weaker than ours ; however, it seems that these methods have not yet been developed so far as to get the central limit theorem with a rate of convergence and the local central limit theorem of this paper. See Section 3 below for more details.

2 Statements of results

For $g \in G$, we set

$$c(g) = \sup \left\{ \frac{d(gx, gy)}{d(x, y)} : x, y \in M, x \neq y \right\},$$

by assumption $c(g) < +\infty$.

For $n \in \mathbb{N}^*$ we denote by π^{*n} the distribution of R_n . We choose a fixed point x_0 in M . For $\eta \geq 1$ and $n \in \mathbb{N}^*$, we define the integrals :

$$\mathcal{M}_\eta = \int_G \left(1 + c(g) + d(gx_0, x_0) \right)^\eta d\pi(g)$$

$$\mathcal{M}'_\eta = \int_G c(g) \left(1 + c(g) + d(gx_0, x_0) \right)^{\eta-1} d\pi(g)$$

$$\mathcal{C}_\eta^{(n)} = \int_G c(g) \max\{c(g), 1\}^{\eta-1} d\pi^{*n}(g).$$

Notice that, since $c(\cdot)$ is submultiplicative, $\mathcal{M}'_\eta < +\infty$ implies $\mathcal{C}_\eta^{(n)} < +\infty$.

The statements below will appeal, on the one hand to the moment conditions $\mathcal{M}_\eta < +\infty$ and $\mathcal{M}'_{\eta'} < +\infty$, on the other hand to the average contractivity condition $\mathcal{C}_{\eta'}^{(n)} < 1$, for a suitable choice of $\eta, \eta' \geq 1$.

We consider a real valued measurable function ξ on $G \times M$ satisfying :

Condition (RS). *There exist $r, s \in \mathbb{R}_+$ and measurable non-negative functions R, S on G such that, for all $x, y \in M$ and $g \in G$,*

$$\begin{aligned} |\xi(g, x)| &\leq R(g)(1 + d(x, x_0))^r \\ |\xi(g, x) - \xi(g, y)| &\leq S(g) d(x, y) \left(1 + d(x, x_0) + d(y, x_0)\right)^s. \end{aligned}$$

Observe that, if the second condition in (RS) holds, then the first one is also valid with $r = s + 1$ and $R(g) = |\xi(g, x_0)| + S(g)$. However, it is worth noticing that this condition may be verified for a smaller exponent r ; this is the case, for example, when ξ is bounded. This remark also shows that, without a significant loss of generality, we could add to condition (RS) the inequality $r \leq s + 1$; yet, we notice that, when r increases, $R(g)$ decreases. The case $s = 0$ and $r = 1$ corresponds to functions ξ such that $\xi(g, \cdot)$ is Lipschitz for all $g \in G$. At last, notice that, if $\alpha \in]0, 1]$, then $d(\cdot, \cdot)^\alpha$ is a distance on M ; consequently Condition (RS) involves the case of functions ξ such that $\xi(g, \cdot)$ is locally α -Hölder for all $g \in G$.

As in the introduction, we denote by Z a r.v. in M defined on (Ω, \mathcal{F}, P) , and independent of the r.v. $Y_n, n \geq 1$. We set

$$S_n^Z = \sum_{k=1}^n \xi(Y_k, Z_{k-1}), \quad n \geq 1.$$

We now state central limit theorems for the sequence $(S_n^Z)_n$; more precise results concerning the behaviour of the sequence $(R_n Z, S_n^Z)_n$ are given in Section 9.

A preliminary to all these statements is the existence of a probability distribution on M which is preserved by the action of π . More precisely, the action on M of the sequence of random mappings $(R_n)_{n \geq 0}$ defines a Markov chain : for $y_0 \in M$, the sequence $(R_n y_0)_{n \geq 0}$ is Markov with state space M , initial distribution δ_{y_0} , and transition probability P defined by

$$y \in M, \quad B \in \mathcal{M}, \quad P(y, B) = \int_G 1_B(gy) d\pi(g).$$

Theorem I (invariant probability measure). *Assume that there exist $\gamma \geq 0$ and an integer $n_0 \geq 1$ such that $\mathcal{M}_{\gamma+1} < +\infty$ and $\mathcal{C}_{\gamma+1}^{(n_0)} < 1$.*

Then there exists on (M, \mathcal{M}) a unique P -invariant probability distribution ν . Moreover we have

$$\int_M d(x, x_0)^{\gamma+1} d\nu(x) < +\infty,$$

and the geometric ergodicity holds in the Prohorov distance d_P . Namely, there exist positive real numbers C and $\kappa_0 < 1$, such that, for any probability distribution μ on M satisfying $\mu(d(\cdot, x_0)) < +\infty$, and all $n \geq 1$,

$$d_P(\mu P^n, \nu) \leq C \kappa_0^{n/2}.$$

It must be noted that such an ergodicity result holds under much weaker hypotheses, see the survey of Diaconis and Freedman (1999) and a recent result in Bhattacharya and Majumdar (2002). In fact, the above statement is just the one which fits the general hypotheses of the paper.

In the sequel our hypotheses will involve a parameter $\gamma_0 > 0$ and

Condition $\mathcal{H}(\gamma_0)$:

$$\mathcal{M}_{\gamma_0+1} < +\infty, \quad \mathcal{M}'_{2\gamma_0+1} < +\infty$$

there exists $n_0 \in \mathbb{N}^*$ such that $\mathcal{C}_{2\gamma_0+1}^{(n_0)} < 1$.

Since $\mathcal{C}_{\gamma_0+1}^{(n_0)} \leq \mathcal{C}_{2\gamma_0+1}^{(n_0)}$, if the above conditions hold, then the P -invariant distribution ν , whose existence is ensured by Theorem I, is such that $\nu(d(\cdot, x_0)^{\gamma_0+1}) < +\infty$; consequently, if the number r and the function R in Condition (RS) verify $r \leq \frac{\gamma_0}{2} + \frac{1}{2}$ and $\int_G R(g)^2 d\pi(g) < +\infty$, we have

$$\int_M \int_G \xi(g, x)^2 d\pi(g) d\nu(x) < +\infty.$$

From now on we shall assume that

$$m = \int_M \int_G \xi(g, x) d\pi(g) d\nu(x) = 0.$$

This causes no loss of generality since it is always possible to replace ξ by $\xi - m$. Otherwise, we shall keep in mind that, if Z has the P -invariant distribution ν , then we have $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$. However, unless otherwise stated, in the sequel Z is not supposed to be ν -distributed.

At last, we define for $g \in G$,

$$\tilde{\delta}(g) = 1 + c(g) + d(gx_0, x_0),$$

and for $\tau > 0$ and positive real valued measurable functions U, V on G , we set

$$\mathcal{J}^\tau(U, V) = \int_G U(g) c(g) \tilde{\delta}(g)^{2\tau} d\pi(g) + \int_G V(g) \tilde{\delta}(g)^{\tau+1} d\pi(g),$$

or more shortly $\mathcal{J}^\tau(U, V) = \pi(Uc\tilde{\delta}^{2\tau}) + \pi(V\tilde{\delta}^{\tau+1})$.

Theorem A (central limit).

Assume $\mathcal{H}(\gamma_0)$ with $\gamma_0 > r + \max\{r, s + 1\}$ and that

$$\int_G R^2 d\pi < +\infty, \quad \mathcal{J}^{\gamma_0-r}(R, R + S) < +\infty.$$

Then there exists a real number $\sigma^2 \geq 0$ such that, under the condition $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$, the sequence $(\frac{S_n^Z}{\sqrt{n}})_{n \geq 1}$ converges in distribution to a $\mathcal{N}(0, \sigma^2)$ -distributed r.v.

As already mentioned, this statement is not the best known one : using our spectral method, it is a stage to the two following results.

Theorem B (central limit with a rate of convergence).

Assume $\mathcal{H}(\gamma_0)$ with $\gamma_0 > 3r + \max\{r, s + 1\}$ and that

$$\int_G R^3 d\pi < +\infty, \quad \mathcal{J}^{\gamma_0-r}(R, R+S) + \mathcal{J}^{\gamma_0-2r}(R^2, (R+S)R) < +\infty.$$

Then, if $\sigma^2 > 0$, there exists a constant C such that, when Z verifies $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$, we have, for all $n \geq 1$,

$$\sup_{u \in \mathbb{R}} \left| \mathbb{P} \left[S_n^Z \leq u\sigma\sqrt{n} \right] - \mathcal{N}(0, 1)(\cdot - \infty, u) \right| \leq C \frac{1 + \mathbb{E}[d(Z, x_0)^{\gamma_0+1}]}{\sqrt{n}}.$$

We denote by \mathcal{L} the Lebesgue's measure on \mathbb{R} . Furthermore, a complex valued function on M is said to be locally Lipschitz if it is Lipschitz on every compact subset of M .

Theorem C (local central limit).

Assume that the conditions of Theorem A hold, and that ξ verifies the non-arithmeticity condition : there is no $t \in \mathbb{R}$, $t \neq 0$, no $\lambda \in \mathbb{C}$, $|\lambda| = 1$, no bounded locally Lipschitz function w on M with non-zero constant modulus on the support Σ_ν of ν , such that we have, for all $x \in \Sigma_\nu$ and all $n \geq 1$,

$$e^{itS_n^x} w(R_n x) = \lambda^n w(x) \quad \mathbb{P} - a.s.$$

Then, if $\sigma^2 > 0$, and if Z is such that $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$, we have, for every continuous function h on \mathbb{R} such that $\lim_{|u| \rightarrow +\infty} u^2 h(u) = 0$,

$$\lim_n \sigma \sqrt{2\pi n} \mathbb{E} \left[h(S_n^Z) \right] = \mathcal{L}(h).$$

We end with a result which gives a criterion for $\sigma^2 > 0$ and defines σ^2 asymptotically.

Theorem S. Assume $\mathcal{H}(\gamma_0)$ with $\gamma_0 > 2r + s + 1$ and that

$$\mathcal{J}^{\gamma_0-r}(R, R+S) + \mathcal{J}^{\gamma_0-2r}(R^2, (R+S)R) < +\infty.$$

(i) If $\sigma^2 = 0$, then there exists a real valued locally Lipschitz function $\tilde{\xi}_1$ on M satisfying $\nu(\tilde{\xi}_1^2) < +\infty$, and such that we have, with Z distributed according to ν ,

$$\xi(Y_1, Z) = \tilde{\xi}_1(Z) - \tilde{\xi}_1(Y_1 Z) \quad \mathbb{P} - a.s.$$

(ii) If the distribution of Z verifies $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$, then

$$\sigma^2 = \lim_n \frac{1}{n} \mathbb{E}[(S_n^Z)^2].$$

It will be seen later on, Theorem C' and S' (§ 9), that the functions w and $\tilde{\xi}_1$ in the two last statements must not be merely locally Lipschitz, they must belong to certain spaces to be defined in the sequel.

In the following section, we show how these theorems apply to some cases of interest. This being done, the rest of the paper is devoted to the proofs ; the reader will find in Subsection 4.2 a brief outline of the subsequent work.

3 Applications.

3.1 Sequences of type $\chi(Z_n)_n$

Let χ be a real valued locally Lipschitz function on M , and suppose that there exist $C, s \in \mathbb{R}_+$ such that, for all $x, y \in M$,

$$|\chi(x) - \chi(y)| \leq Cd(x, y)(1 + d(x, x_0) + d(y, x_0))^s.$$

Using martingale methods, it is proved that the central limit theorem for $(\chi(Z_n))_n$ holds for any initial distribution under the moment condition $\int_G d(gx_0, x_0)^{4(s+1)} d\pi(g) < +\infty$ and the contraction property $\int_G c(g)^{4(s+1)} d\pi(g) < 1$, See Dufflo (1997). By means of similar techniques, it is established in Benda (1998), that, when $s = 0$, the same result is valid under the weaker hypotheses $\int_G c(g)^2 d\pi(g) < 1$ and $\int_G d(gx_0, x_0)^2 d\pi(g) < +\infty$. Considering the stationary chain with initial probability ν , Wu and Woodroffe (2000) have established a central limit theorem for functions χ which are not Lipschitz and not even continuous.

Let us now apply the results of the preceding section : we set $\xi(g, x) = \chi(x)$. The moment hypotheses of Theorem A are the same as those of Theorem C, so that it can be seen from Theorem 3.1 below that they are stronger than the ones previously stated. However, to our knowledge, Theorem B and C are new. They can be stated as follows. Recall that $\tilde{\delta}(g) = 1 + c(g) + d(gx_0, x_0)$.

Theorem 3.1. *Suppose that there exist $\epsilon > 0$ and integers $n_0 \geq 1, k \geq 0$ such that*

$$\pi\left(\tilde{\delta}^{k(s+1)+1+\frac{\epsilon}{2}} + c\tilde{\delta}^{2k(s+1)+\epsilon}\right) < +\infty, \quad \text{and} \quad \pi^{*n_0}\left(c \max\{1, c\}^{2k(s+1)+\epsilon}\right) < 1,$$

and assume that $\nu(\chi) = 0$, where ν is the P -invariant probability measure.

If k takes the values 4 and 3 respectively, then the assertions of Theorems B and S respectively

apply to $S_n^Z = \sum_{k=1}^n \chi(Z_{k-1})$.

Moreover, if χ is non-arithmetic and if the above integral conditions are satisfied for $k = 2$, then the assertion of Theorem C holds.

Proof. The function ξ on $G \times M$ defined by $\xi(g, y) = \chi(y)$ verifies Condition (RS) with the exponents $r = s + 1$ and s associated with constant functions R and S . These have moments of all orders. Consequently the moment conditions of theorems B, S, and C reduce to $\mathcal{H}(\gamma_0)$ with $\gamma_0 = k(s + 1) + \frac{\varepsilon}{2}$; this gives the desired results. \square

Let us point out that Pollicott (2001) has stated a central limit theorem with a rate of convergence and a large deviations theorem in the case where the support of the probability measure π is finite. However this study is based on the assertion without proof that, on a suitable space of Lipschitz functions, the Fourier kernels $P(t)$ (See § 4) are analytic perturbed operators of P . Also notice that, if it is proved that the stationary chain with initial probability distribution ν is strongly mixing and Harris recurrent, then we can apply Bolthausen (1982) to obtain a central limit theorem with a $n^{-1/2}$ rate of convergence. However, on one hand this requires some additional hypotheses on π (see Meyn and Tweedie (1993) p. 140 for a sufficient condition in the context of the following section), on the other hand this only covers the stationary case.

3.2 Generalized autoregressive processes

Denote by G the semigroup of all affine mappings of $M = \mathbb{R}^q$, $q \geq 1$. An element $g \in G$ is identified with a couple $(a(g), b(g))$, where $a(g)$ is an endomorphism of \mathbb{R}^q and $b(g)$ is a vector in \mathbb{R}^q . For $y \in M$, we set $gy = a(g)y + b(g)$. The associated generalized autoregressive process $(Z_n)_{n \geq 0}$ is then defined by

$$Z_0 = Z, \quad Z_{n+1} = a(Y_{n+1})Z_n + b(Y_{n+1}), \quad n \geq 0.$$

Let ξ be a function from $G \times \mathbb{R}^q$ to \mathbb{R} , and suppose that there exist a norm $\|\cdot\|$ on \mathbb{R}^q , $\alpha \in]0, 1]$, $r, s \in \mathbb{R}_+$ and non-negative measurable functions R and S on G such that, for all $g \in G$ and $x, y \in \mathbb{R}^q$, we have

$$|\xi(g, x)| \leq R(g)(1 + \|x\|)^{\alpha r}, \quad |\xi(g, x) - \xi(g, y)| \leq S(g)\|x - y\|^\alpha(1 + \|x\| + \|y\|)^{\alpha s}.$$

For instance, these properties hold with $\alpha = 1$ when ξ is a polynomial function of the entries of the matrix representing $a(g)$ and of the coordinates of the vectors $b(g)$ and x .

Let us consider the distance d defined on \mathbb{R}^q by $d(x, y) = \|x - y\|^\alpha$, and choose $x_0 = 0 \in \mathbb{R}^q$. We have $c(g) = \|a(g)\|^\alpha$ and $d(gx_0, x_0) = \|b(g)\|^\alpha$. Then the statements B-C-S apply straightforwardly. To compare with former results, let us rewrite Theorem B. Let $\tilde{\delta}(g) = (1 + \|a(g)\| + \|b(g)\|)^\alpha$, then

Theorem 3.2. *The hypotheses in central limit theorem with a rate of convergence (Th. B) are satisfied if there exist $\gamma_0 > 3r + \max\{r, s + 1\}$ and $n_0 \in \mathbb{N}^*$ such that*

$$\pi\left(\tilde{\delta}^{\gamma_0+1} + \|a\|^\alpha \tilde{\delta}^{2\gamma_0}\right) < +\infty, \quad \text{and} \quad \pi^{*n_0}\left(\|a\|^\alpha \max\{1, \|a\|\}^{2\gamma_0\alpha}\right) < 1,$$

and when the functions $R(\cdot)$ and $S(\cdot)$ satisfy the moment conditions

$$\int_G R^3 d\pi < +\infty, \quad \mathcal{J}^{\gamma_0-r}(R, R+S) + \mathcal{J}^{\gamma_0-2r}(R^2, (R+S)R) < +\infty.$$

In this context, convergence rates in the central limit theorem have already been established by Milhaud and Raugi (1989), and by Cuny (2003). The spectral method used in Milhaud and Raugi (1989) is, in substance, similar to the one developed here, but it appeals to the standard perturbation theorem : for this reason (See § 6.1), the following conditions on $a(\cdot)$ and $b(\cdot)$ are required : $\|a(\cdot)\| < 1$ π -p.s, and there exist real numbers $\rho > 0$ and $\beta \in]0, 1]$ such that we have the exponential moment condition

$$\int_G e^{\rho \|b(g)\|^\beta} (R(g) + S(g))^5 (1 - \|a(g)\|^\beta)^{-\frac{5\rho}{\beta}(1+\max\{r,s+1\})} d\pi(g) < +\infty.$$

The hypotheses on both $a(\cdot)$ and $b(\cdot)$ are significantly less restrictive in Theorem 3.2. The study in Cuny (2003) is based on martingale methods. The contraction condition is the same as in Milhaud and Raugi (1989), and it is supposed that, for all $\ell \in \mathbb{N}$, $\int \|b(g)\|^\ell d\pi(g) < +\infty$. Under these conditions, for functions ξ which are not necessarily Hölder of the variable x , it is proved that the rate of convergence in the central limit theorem is n^{-p} for every $p < 1/2$.

3.3 Products of positive random matrices

Let G be the semi-group of $q \times q$ matrices with non-negative entries which are allowable, namely every row and every column contains a strictly positive element, and denote by G° the ideal of G composed of matrices with strictly positive entries.

For $g \in G$ and $w \in \mathbb{R}^q$, we denote by $g(w)$ the image of w under g ; the cone

$$C = \{w : w = (w_1, \dots, w_q) \in \mathbb{R}^q, w_k > 0, k = 1, \dots, q\}$$

is invariant under all $g \in G$. Define M to be the intersection of the hyperplane $\{w : w \in \mathbb{R}^q, \sum_{k=1}^q w_k = 1\}$ of \mathbb{R}^q with C .

The linear space \mathbb{R}^q is endowed with the norm $\|\cdot\|$ defined by

$$w = (w_1, \dots, w_q) \in \mathbb{R}^q, \quad \|w\| = \sum_{k=1}^q |w_k|,$$

and, for each $g \in G$, we set

$$\|g\| = \sup\{\|g(y)\| : y \in M\}, \quad v(g) = \inf\{\|g(y)\| : y \in M\}.$$

The semi-group G being equipped with its Borel σ -field \mathcal{G} , we consider a probability distribution π on G for which there exists an integer n_0 such that the support of the r.v. R_{n_0} contains a matrix of G° . Denote by g^* the adjoint of g , it is shown in Hennion (1997) that, if

$$\int_G \left(|\ln \|g^*\| | + |\ln v(g^*)| \right)^2 d\pi(g) < +\infty,$$

then there exists $\gamma_1 \in \mathbb{R}$ such that, for $y \in M$, the sequence $\left(\frac{1}{\sqrt{n}} (\ln \|R_n(y)\| - n\gamma_1) \right)_{n \geq 1}$ converges to the $\mathcal{N}(0, \sigma^2)$ distribution, moreover the case $\sigma^2 = 0$ is investigated. Using

Theorems B and C, it is possible to state a central limit theorem with a rate of convergence and a local central limit theorem. Notice that similar theorems have already been given in Hennion and Hervé (2001) Section X.5. but under more restrictive moment hypotheses.

To see how this case enters the present frame, we set for $g \in G$ and $y \in M$

$$gy = \frac{g(y)}{\|g(y)\|}, \quad a(g, y) = \ln \|g(y)\|,$$

It is easy to check that the first formula defines an action of G on M , while the function defined by the second one verifies the property of additive cocycle associated with this action :

$$a(gg', y) = a(g, g'y) + a(g', y), \quad g, g' \in G, \quad y \in M.$$

Consequently, setting $\xi(g, y) = a(g, y) - \gamma_1$, for $(g, y) \in G \times M$, we can write, for $y \in M$,

$$\ln \|R_n(y)\| - n\gamma_1 = \sum_{k=1}^n \xi(Y_k, R_{k-1}y).$$

Furthermore, when M is endowed with a suitable metric d_H called the Hilbert metric, see Bapat and Raghavan (1997), every $g \in G$ is Lipschitz with constant $c(g) \leq 1$, and we have $c(g) < 1$ if and only if all entries of g are strictly positive. Therefore, if the support of R_{n_0} contains such a matrix g , we have $\mathcal{C}_\eta^{(n_0)} = \int_G c(g) d\pi^{*n_0}(g) < 1$ for all $\eta \geq 1$.

For $\eta \geq 0$, set

$$\mathcal{L}^\eta = \int_G \left(|\ln \|g\|| + |\ln v(g)| + |\ln v(g^*)| \right)^\eta d\pi(g).$$

Theorem 3.3. *Suppose that there exists an integer n_0 such that the support of the r.v. R_{n_0} contains a matrix of G° , and let $\epsilon > 0$.*

(i) *Assume $\mathcal{L}^{4+\epsilon} < +\infty$, then, if $\sigma^2 > 0$, there exists a non-negative constant C such that, in case the r.v. Z of M verifies $\mathbb{E}[d_H(Z, x_0)^{2+\frac{\epsilon}{2}}] < +\infty$, we have, for all $n \geq 1$,*

$$\sup_{u \in \mathbb{R}} \left| \mathbb{P} \left[\ln \|R_n(Z)\| - n\gamma_1 \leq u\sigma\sqrt{n} \right] - \mathcal{N}(0, 1)(-\infty, u) \right| \leq C \frac{1 + \mathbb{E}[d_H(Z, x_0)^{2+\frac{\epsilon}{2}}]}{\sqrt{n}}.$$

(ii) *Assume $\mathcal{L}^{3+\epsilon} < +\infty$, $\sigma^2 > 0$, and that the support of R_{n_0} contains two matrices $g_1, g_2 \in G^\circ$ whose spectral radii ρ_1, ρ_2 verify $\ln \frac{\rho_2}{\rho_1} \notin \mathbb{Q}$. Then, if $\mathbb{E}[d_H(Z, x_0)^{2+\frac{\epsilon}{2}}] < +\infty$, we have, for any real valued continuous function h on \mathbb{R} such that $\lim_{|u| \rightarrow +\infty} u^2 h(u) = 0$,*

$$\lim_n \sigma \sqrt{2\pi n} \mathbb{E} \left[h(\ln \|R_n(Z)\| - n\gamma_1) \right] = \mathcal{L}(h).$$

Proof. First notice that the number $\|g^*\|$ associated to any endomorphism g of \mathbb{R}^q defines a new norm which is equivalent to the one already considered. Consequently there exists a constant C such that, for every $g \in G$, we have $|\ln \|g^*\|| \leq C + |\ln \|g\||$.

We denote by $(e_k)_{k=1}^q$ and by $\langle \cdot, \cdot \rangle$ the canonical basis and scalar product on \mathbb{R}^q . If $y, y' \in M$, we set

$$m_H(y, y') = \min \left\{ \frac{\langle y, e_k \rangle}{\langle y', e_k \rangle} : k = 1, \dots, q \right\}, \quad d_H(y, y') = -\ln(m_H(y, y')m_H(y', y)),$$

d_H is the Hilbert distance on M , see Bapat and Raghavan (1997). The space (M, d_H) is not compact, but each closed ball in it is compact. Set $x_0 = (1/q, \dots, 1/q) \in M$, we have

$$d(gx_0, x_0) = \ln \frac{\max_i \langle gx_0, e_i \rangle}{\min_j \langle gx_0, e_j \rangle} = \ln \frac{\max_i \|g^* e_i\|}{\min_j \|g^* e_j\|} = \ln \frac{\|g^*\|}{v(g^*)} \leq C + |\ln \|g\|| + |\ln v(g^*)|.$$

The function $\xi(g, \cdot)$ is bounded by $|\gamma_1| + |\ln \|g\|| + |\ln v(g)|$. From $\|gy\| \geq m_H(y, y') \|gy'\|$ we deduce that $S(g) = 1$ and $s = 0$, see Hennion (1997) Lemma 5.3. Therefore Condition (RS) is verified with $R(g) = 2(|\ln \|g\|| + |\ln v(g)|)$, $r = 0$, and $S(g) = 1$, $s = 0$.

The above estimations prove that the required moment conditions of theorems B and C hold if we have respectively $\mathcal{L}^{4+\epsilon} < +\infty$ and $\mathcal{L}^{3+\epsilon} < +\infty$.

It remains to prove that the additional hypothesis in (ii) implies the non-arithmeticity of ξ . Let $k = 1, 2$. It follows from the Perron-Frobenius' Theorem that $\rho_k > 0$ and that, for all $\ell \geq 1$, we have $g_k^\ell = \rho_k^\ell (p_k + h_k^\ell)$, where $p_k \in G^\circ$ and the endomorphism h_k of \mathbb{R}^q has a spectral radius < 1 . Consequently, for any $x \in M$, we have $\ln \|g_k^\ell x\| = \ell \ln \rho_k + r_{k,\ell}(x)$, with $\lim_\ell r_{k,\ell}(x) = \ln \|p_k(x)\|$. Suppose that there exist $t \in \mathbb{R}$, $t \neq 0$, $\lambda \in \mathbb{C}$, $|\lambda| = 1$, and a bounded locally Lipschitz function w on M which has a non-zero constant modulus on the support Σ_ν of ν , and such that we have, for all $x \in \Sigma_\nu$ and all $n \geq 1$,

$$\lambda^n w(x) = e^{itS_n^x} w(R_n x) = e^{it(\ln \|R_n(x)\| - n\gamma_1)} w(R_n x) \quad P - a.s.$$

From the continuity of the functions used in the two members, we deduce that, for any $\ell \geq 1$ and $x \in \Sigma_\mu$, we have

$$e^{it(\ln \|g_k^\ell(x)\| - n_0 \ell \gamma_1)} w(g_k^\ell x) = \lambda^{n_0 \ell} w(x).$$

It follows that

$$e^{it\ell \ln \frac{\rho_2}{\rho_1}} = \frac{w(g_1^\ell x)}{w(g_2^\ell x)} e^{it(r_{1,n_0 \ell}(x) - r_{2,n_0 \ell}(x))}.$$

The second member converges when $\ell \rightarrow +\infty$, while the countable set of complex numbers defined by the first one is dense in $\{z : z \in \mathbb{C}, |z| = 1\}$. This contradiction completes the proof. \square

4 Preliminaries

4.1 P -invariant probability measure (proof of Theorem I)

Here the hypotheses are those of Theorem I : $\mathcal{M}_{\gamma+1} < +\infty$, $\mathcal{C}_{\gamma+1}^{(n_0)} < 1$ ($\gamma \geq 0$, $n_0 \in \mathbb{N}^*$). For $\lambda \in]0, 1]$, $x \in M$, and $g \in G$, we set

$$p_\lambda(x) = 1 + \lambda d(x, x_0), \quad \delta_\lambda(g) = \max\{c(g), 1\} + \lambda d(gx_0, x_0), \quad \text{and} \quad \tilde{\delta}(g) = 1 + c(g) + d(gx_0, x_0).$$

Lemma 4.1. *We have for all $g \in G$ and $0 \leq \lambda \leq 1$*

$$\sup_{x \in M} \frac{p_\lambda(gx)}{p_\lambda(x)} \leq \delta_\lambda(g) \leq \tilde{\delta}(g).$$

The functions $c(\cdot)$ and $\tilde{\delta}(\cdot)$ are submultiplicative.

Proof. Let $x \in M$ and $g \in G$, then

$$\begin{aligned} \frac{p_\lambda(gx)}{p_\lambda(x)} &= \frac{1 + \lambda d(gx, x_0) - \lambda d(gx_0, x_0)}{1 + \lambda d(x, x_0)} + \frac{\lambda d(gx_0, x_0)}{1 + \lambda d(x, x_0)} \\ &\leq \frac{1 + \lambda d(gx, gx_0)}{1 + \lambda d(x, x_0)} + \lambda d(gx_0, x_0) \leq \frac{1 + \lambda c(g)d(x, x_0)}{1 + \lambda d(x, x_0)} + \lambda d(gx_0, x_0) \\ &\leq \max\{1, c(g)\} + \lambda d(gx_0, x_0). \end{aligned}$$

The fact that $c(\cdot)$ is submultiplicative is obvious. Finally for $h, g \in G$, we get

$$\begin{aligned} \tilde{\delta}(hg) &\leq 1 + c(h)c(g) + [d(hgx_0, x_0) - d(hx_0, x_0)] + d(hx_0, x_0) \\ &\leq 1 + c(h)c(g) + c(h)d(gx_0, x_0) + d(hx_0, x_0) \leq \tilde{\delta}(h)\tilde{\delta}(g). \end{aligned}$$

□

Recall that, for $n \in \mathbb{N}^*$, π^{*n} denotes the law of R_n .

Lemma 4.2. *Let $\phi_\lambda(x) = d(x, x_0)p_\lambda(x)^\gamma$ for $\lambda \in [0, 1]$. Then*

- (a) *For all $n \geq 1$ and $x \in M$, we have $P^n \phi_\lambda(x) < +\infty$.*
- (b) *For $\lambda_0 \in]0, 1]$ small enough, we have $\int_G c(g)\delta_{\lambda_0}(g)^\gamma d\pi^{*n_0}(g) < 1$.*
- (c) *There exist constants $\varepsilon \in]0, 1[$, $C \in \mathbb{R}_+$ such that*

$$P^{n_0} \phi_{\lambda_0} \leq C + \varepsilon \phi_{\lambda_0}.$$

Proof. (a) For $n \geq 1$ and $x \in M$, we have

$$\begin{aligned} P^n \phi_\lambda(x) &= \int_G d(gx, x_0)p_\lambda(gx)^\gamma d\pi^{*n}(g) \\ &\leq \int_G d(gx_0, x_0)p_\lambda(gx)^\gamma d\pi^{*n}(g) + \int_G [d(gx, x_0) - d(gx_0, x_0)]p_\lambda(gx)^\gamma d\pi^{*n}(g) \\ &\leq p_\lambda(x)^\gamma \int_G d(gx_0, x_0)\delta_\lambda(g)^\gamma d\pi^{*n}(g) + d(x, x_0)p_\lambda(x)^\gamma \int_G c(g)\delta_\lambda(g)^\gamma d\pi^{*n}(g). \end{aligned}$$

The functions in the two integrals above are dominated by $\tilde{\delta}(\cdot)^{\gamma+1}$. Since this function is submultiplicative and π -integrable, Fubini's theorem ensures that these integrals are finite. Thus $P^n \phi_\lambda(x) < +\infty$.

Since $c\delta_{\lambda_0}^\gamma \leq \tilde{\delta}^{\gamma+1}$ and $\tilde{\delta}$ is submultiplicative, Assertion (b) is a direct consequence of Hypotheses and Lebesgue's theorem.

Set $\varepsilon' = \int_G c(g)\delta_{\lambda_0}(g)^\gamma d\pi^{*n_0}(g)$. From the above inequality applied with $\lambda = \lambda_0$ and $n = n_0$, there exists a constant D_0 such that

$$P^{n_0}\phi_{\lambda_0} \leq D_0 p_{\lambda_0}^\gamma + \varepsilon'\phi_{\lambda_0}.$$

Using continuity and $\lim_{d(x,x_0) \rightarrow +\infty} \frac{p_{\lambda_0}(x)^\gamma}{\phi_{\lambda_0}(x)} = 0$, we see that there exists a constant C such that $D_0 p_{\lambda_0}^\gamma \leq C + \frac{1-\varepsilon'}{2}\phi_{\lambda_0}$. Hence $P^{n_0}\phi_{\lambda_0} \leq C + \varepsilon\phi_{\lambda_0}$ with $\varepsilon = \frac{1+\varepsilon'}{2}$. \square

Now let us prove Theorem I. For convenience we set $\phi = \phi_{\lambda_0}$. By induction and Lemma 4.2, we obtain, for every $q \geq 1$, $P^{qn_0}\phi \leq \varepsilon^q\phi + C(1 + \varepsilon + \dots + \varepsilon^{q-1}) \leq \phi + \frac{C}{1-\varepsilon}$. Let $n \in \mathbb{N}^*$. Writing $n = qn_0 + r$ with $r \in \{0, \dots, n_0 - 1\}$ and setting $E = \max\{P^k\phi(x_0), k = 0, \dots, n_0 - 1\}$, we get $P^n\phi(x_0) \leq E + \frac{C}{1-\varepsilon}$. Therefore the sequence $(P^n\phi(x_0))_n$ is bounded by a constant, say K . For $n \geq 1$, let ν_n be the probability measure on (M, \mathcal{M}) defined by

$$B \in \mathcal{M}, \quad \nu_n(B) = \frac{1}{n} \sum_{k=0}^{n-1} (P^k 1_B)(x_0).$$

Observe that, for each $n \geq 1$, we have $\nu_n(\phi) \leq K$. Since $\lim_{d(x,x_0) \rightarrow +\infty} \phi(x) = \infty$, the subset $[\phi \leq \alpha]$ is compact for each $\alpha > 0$. The Markov's inequality implies that, for all $n \geq 1$, we have $\nu_n([\phi > \alpha]) \leq \frac{\nu_n(\phi)}{\alpha} \leq \frac{K}{\alpha}$, so that the sequence $(\nu_n)_n$ is tight. Therefore we can select a subsequence $(\nu_{n_k})_k$ converging to a probability measure ν . It is clear that ν is P -invariant.

For $p \in \mathbb{N}^*$, set $\phi_p(\cdot) = \min(\phi(\cdot), p)$. For $k \geq 0$ and $p \geq 0$, we have $\nu_{n_k}(\phi_p) \leq \nu_{n_k}(\phi) \leq K$, consequently, for all $p \geq 0$, $\lim_k \nu_{n_k}(\phi_p) = \nu(\phi_p) \leq K$. The monotone convergence theorem gives $\nu(\phi) < +\infty$, that is $\nu(d(\cdot, x_0)^{\gamma+1}) < +\infty$.

Now let us prove that ν is the unique P -invariant probability distribution. First observe that, since $\mathbb{E}[\ln c(R_{n_0})] \leq \ln \mathbb{E}[c(R_{n_0})] = \ln \mathcal{C}_1^{(n_0)} < 0$, the law of large numbers asserts that $\limsup_q c(R_{qn_0})^{\frac{1}{q}} \leq \lim_q (\prod_{\ell=1}^q c(Y_{\ell n_0} \dots Y_{(\ell-1)n_0+1}))^{1/q} < 1$ on a set Ω_1 such that $\mathbb{P}(\Omega_1) = 1$. For $x, y \in M$ and $q \geq 1$, we can write $d(R_{qn_0}x, R_{qn_0}y) \leq c(R_{qn_0})d(x, y)$, so that $\lim_q d(R_{qn_0}x, R_{qn_0}y) = 0$ on Ω_1 . Let ν' be a P -invariant probability distribution on M . For each bounded continuous function f on M , we have

$$\nu'(f) - \nu(f) = \int_M E[f(R_{qn_0}x) - f(R_{qn_0}y)] d\nu'(x) d\nu(y),$$

passing to the limit, we get $\nu'(f) - \nu(f) = 0$. We conclude that $\nu' = \nu$.

It remains to establish the geometric ergodicity in the Prohorov distance d_P . Let f be a bounded uniformly lipschitz function on M . Then, for all $x, y \in M$ and $n \geq 1$, we have

$$\begin{aligned} |P^n f(x) - P^n f(y)| &\leq \int_G |f(gx) - f(gy)| d\pi^{*n}(g) \leq m_0(f)d(x, y) \int_G \frac{d(gx, gy)}{d(x, y)} d\pi^{*n}(g) \\ &\leq m_0(f)d(x, y) \mathcal{C}_1^{(n)} \end{aligned}$$

where $m_0(f) = \sup\{\frac{|f(x) - f(y)|}{d(x, y)}, x, y \in M, x \neq y\}$. Let μ be the law of Z , and assume that $\phi_0(\cdot) = d(\cdot, x_0)$ is μ -integrable. By integrating the previous inequality with respect to both $d\nu(x)$ and $d\mu(y)$, it follows that $|\nu(f) - \mu P^n(f)| \leq \mathcal{C}_1^{(n)} m_0(f)(\nu(\phi_0) + \mu(\phi_0))$. This bound proves that $\nu - \mu P^n$ is a continuous linear functional on the space of all bounded uniformly lipschitz functions on M endowed with its canonical norm. Moreover we have $\|\nu - \mu P^n\| \leq C' \mathcal{C}_1^{(n)}$ with $C' = \nu(\phi_0) + \mu(\phi_0)$. Writting $n = qn_0 + r$ with $r \in \{0, \dots, n_0 - 1\}$ and using the fact that $c(\cdot)$ is submultiplicative, we easily see that $\mathcal{C}_1^{(n)} \leq C'' (\mathcal{C}_1^{(n_0)})^{\frac{n}{n_0}}$. Since $d_p(\nu, \mu P^n) \leq 2\|\nu - \mu P^n\|^{\frac{1}{2}}$, see Dudley (1989), the last assertion of Theorem I follows with $\kappa_0 = (\mathcal{C}_1^{(n_0)})^{\frac{1}{n_0}}$. \square

4.2 Outlines of the method

As mentionned in Introduction, the main idea of this work consists in applying the method described in Hennion and Hervé (2001) to the function ξ and to the Markov chain $(X_n)_{n \geq 0}$ with the state space $G \times M$ and the transition probability Q defined by

$$(g, y) \in G \times M, \quad B \in \mathcal{G} \times \mathcal{M}, \quad Q((g, y), B) = \int_G 1_B(h, gy) d\pi(h).$$

However, we observed in Chapter X of Hennion and Hervé (2001) devoted to Lipschitz kernels, that, because of the special form of Q , the essential part of the study can be performed with the help of the transition probability P and of the Fourier kernels $P(t)$, $t \in \mathbb{R}$, associated to P and ξ , which are defined, for any bounded measurable function f on M , by

$$y \in M, \quad P(t)f(y) = \int_G e^{it\xi(g, y)} f(gy) d\pi(g).$$

This is due to the fact that, for all functions f as above, we have $Q(f \circ j) = (Pf) \circ j$, where j is the action of G on M . Then, in the sequel, we shall only use the kernels $P(t)$; the next statement indicates that these kernels are sufficient for our purpose.

Basic Lemma. *Let f be a bounded measurable function on M , and denote by μ the distribution of Z . Then we have for $n \geq 1$, $t \in \mathbb{R}$*

$$\mathbb{E}\left[f(R_n Z) e^{itS_n^Z}\right] = \mu(P(t)^n f).$$

Proof. Set $S_0^Z = 0$. For $n \geq 1$, we have

$$\mathbb{E}\left[f(R_n Z) e^{itS_n^Z}\right] = \mathbb{E}\left[f(Y_n R_{n-1} Z) e^{it(S_{n-1}^Z + \xi(Y_n, R_{n-1} Z))}\right].$$

Since (Z, Y_1, \dots, Y_{n-1}) and Y_n are independent r.v., Fubini's theorem gives

$$\begin{aligned} \mathbb{E}\left[f(R_n Z) e^{itS_n^Z}\right] &= \mathbb{E}\left[e^{itS_{n-1}^Z} \int_G f(gR_{n-1} Z) e^{it\xi(g, R_{n-1} Z)} d\pi(g)\right] \\ &= \mathbb{E}\left[e^{itS_{n-1}^Z} (P(t)f)(R_{n-1} Z)\right]. \end{aligned}$$

The desired formula for $n = 1$ holds because the second member equals $\mathbb{E}[P(t)f(Z)] = \mu(P(t)f)$. Suppose now that the stated formula is valid at rank $n - 1$, $n \geq 2$. Then, from the previous relation and the fact that $P(t)f$ is a bounded measurable function on M , we conclude that $\mathbb{E}[f(R_n Z)e^{itS_n^Z}] = \mu(P(t)^{n-1}(P(t)f))$. This completes the proof. \square

Theorems A-B-C-S will be direct consequences of the extensions A'-B'-C'-S' stated in Section 9. The outline of the argumentation is the following. In Section 5, we shall introduce spaces \mathcal{B}_γ , which depend on a real parameter $\gamma > 0$ and are composed of locally Lipschitz functions on M . Three norms denoted by $N_{\infty,\gamma}$, N_γ and $N_{1,\gamma}$, will be defined on \mathcal{B}_γ , it will be proved that they are equivalent, but each of them will be suited to a part of the proof. In this way, in Section 5.3, we shall see that the use of N_γ is convenient to establish that, for suitable γ , P is quasi-compact on \mathcal{B}_γ , and furthermore that the number 1 is the unique peripheral eigenvalue of P . In Section 6 the norms $N_{\infty,\gamma}$ will be helpful for the study of the behaviour of the function $P(t)$ near $t = 0$. For this purpose, it will be worth noticing that, for $\gamma' < \gamma$, $P(t)$ may be viewed as a bounded linear map from $\mathcal{B}_{\gamma'}$ to \mathcal{B}_γ ; indeed the derivative kernels of $P(t)$, which in general do not define bounded endomorphisms of $(\mathcal{B}_\gamma, N_{\infty,\gamma})$, can be considered on the other hand as bounded linear maps from $\mathcal{B}_{\gamma'}$ to \mathcal{B}_γ for suitable $\gamma' < \gamma$; of course this will be a less restricting property because the space \mathcal{B}_γ strictly contains $\mathcal{B}_{\gamma'}$ and is endowed with a weaker norm. In Section 7, the norm $N_{1,\gamma}$ will be an essential tool to apply a perturbation Theorem due to G. Keller and C. Liverani from which it will follow that $P(t)$ are perturbed operators of P for small $|t|$. The interest of this perturbation theorem is that it only requires $P(\cdot)$ to be continuous as a map taking values in the space of bounded linear map from $(\mathcal{B}_\gamma, N_{1,\gamma})$ to $(\mathcal{B}_\gamma, \nu(|\cdot|))$; this is the key point of this study, See § 6.1. In particular this theorem ensures that, for small $|t|$, $P(t)$ has only one dominating simple eigenvalue, $\lambda(t)$, on \mathcal{B}_γ , and we shall establish in Section 8 that the Taylor's expansions for $P(t)$ at $t = 0$ obtained in Section 6 lead to expansions of the eigenelements belonging to $\lambda(t)$. Then, in Section 9, by using the previous preparation and by applying the method described in Hennion and Hervé (2001), we shall be in a position to prove limit theorems. Notice that renewal and large deviations theorems for the sequence $(S_n^Z)_{n \geq 1}$ might be derived from similar techniques.

5 The space \mathcal{B}_γ and quasi-compactness of P

5.1 Conventions and notations

From now on, we fix $\gamma_0 > 0$ and $n_0 \in \mathbb{N}^*$ such that Condition $\mathcal{H}(\gamma_0)$ holds, that is :

$$\mathcal{M}_{\gamma_0+1} = \pi(\tilde{\delta}^{\gamma_0+1}) < +\infty, \quad \mathcal{M}'_{2\gamma_0+1} = \pi(c\tilde{\delta}^{2\gamma_0}) < +\infty, \quad \mathcal{C}_{2\gamma_0+1}^{(n_0)} = \pi^{*n_0}(c \max\{c, 1\}^{2\gamma_0}) < 1.$$

According to the subsequent statements, some additional conditions will be imposed on γ_0 .

Lemma 5.1. *There exists a real number $\lambda_0 \in]0, 1]$ such that*

$$\vartheta_0 = \int_G c(g) \left(\max\{c(g), 1\} + \lambda_0 d(gx_0, x_0) \right)^{2\gamma_0} d\pi^{*n_0}(g) < 1.$$

Proof. Since $c(g) (\max\{c(g), 1\} + \lambda_0 d(gx_0, x_0))^{2\gamma_0} \leq c(g) \tilde{\delta}(g)^{2\gamma_0}$, and the functions c , $\tilde{\delta}$ are submultiplicative, the lemma follows from the two last conditions of $\mathcal{H}(\gamma_0)$ and Lebesgue's theorem. \square

Now we fix a $\lambda_0 \in]0, 1]$ satisfying the previous inequality.

For $x, y \in M$, $g \in G$, we set

- $p(x) = 1 + \lambda_0 d(x, x_0)$
- $\delta(g) = \max\{c(g), 1\} + \lambda_0 d(gx_0, x_0)$.

Notice that $p \leq 1 + d(\cdot, x_0) \leq \frac{1}{\lambda_0} p$, and $\delta \leq \tilde{\delta} \leq \frac{2}{\lambda_0} \delta$. Besides, for $\gamma > 0$, let us write

- $\Delta_\gamma(x, y) = d(x, y) p(x)^\gamma p(y)^\gamma$.

With the help of these elements, we now define the space \mathcal{B}_γ composed of locally Lipschitz functions on M , and we define four equivalent norms on this space. Such spaces, introduced in Le Page (1983), have already been used by several authors in order to prove the quasi-compactness of probability kernels having a contracting property, See Milhaud and Raugi (1989), Peigné (1993) ; a similar statement will be established in Section 5.3.

5.2 Definitions of \mathcal{B}_γ and of the norms $N_{\infty, \gamma}$, $N_{\infty, \gamma, \tilde{\gamma}}$, N_γ and $N_{1, \gamma}$

For $\gamma > 0$, we denote by \mathcal{B}_γ the space of all complex valued locally Lipschitz functions on M such that

$$m_\gamma(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\Delta_\gamma(x, y)}, x, y \in M, x \neq y \right\} < +\infty.$$

The inequality $\Delta_\gamma(x, x_0) = d(x, x_0) p(x)^\gamma \leq \frac{1}{\lambda_0} p(x)^{\gamma+1}$ ensures that, for all $f \in \mathcal{B}_\gamma$, we have $|f(x)| \leq |f(x_0)| + \frac{1}{\lambda_0} m_\gamma(f) p(x)^{\gamma+1}$, thus $\sup_{x \in M} \frac{|f(x)|}{p(x)^{\gamma+1}} < +\infty$. Consequently \mathcal{B}_γ can be equipped with the norm

- $N_{\infty, \gamma}(f) = m_\gamma(f) + |f|_\gamma$, where $|f|_\gamma = \sup \left\{ \frac{|f(x)|}{p(x)^{\gamma+1}}, x \in M \right\}$.

Let $\tilde{\gamma} > \gamma$. As $p^{\tilde{\gamma}+1} \geq p^{\gamma+1}$, we have, for $f \in \mathcal{B}_\gamma$, $|f|_{\tilde{\gamma}} = \sup_{x \in M} \frac{|f(x)|}{p(x)^{\tilde{\gamma}+1}} < +\infty$, we set

- $N_{\infty, \gamma, \tilde{\gamma}}(f) = m_\gamma(f) + |f|_{\tilde{\gamma}}$.

Since $\mathcal{M}_{\gamma_0+1} < +\infty$ and $\mathcal{C}_{\gamma_0+1}^{(n_0)} \leq \mathcal{C}_{2\gamma_0+1}^{(n_0)} < 1$, the P -invariant probability measure, ν , whose existence is ascertained by Theorem I, is such that

$$\int_M d(x, x_0)^{\gamma_0+1} d\nu(x) < +\infty.$$

Therefore, for every $\gamma \in]0, \gamma_0]$, ν integrates $p^{\gamma+1}$, and thus integrates all the functions of \mathcal{B}_γ , so that we can define on \mathcal{B}_γ the following norms

- $N_\gamma(f) = m_\gamma(f) + |\nu(f)|$
- $N_{1,\gamma}(f) = m_\gamma(f) + \nu(|f|)$

Proposition 5.2. *Let γ , $0 < \gamma \leq \gamma_0$. The four norms $N_{\infty,\gamma}$, $N_{\infty,\gamma,\tilde{\gamma}}$, N_γ and $N_{1,\gamma}$ are equivalent on \mathcal{B}_γ . When equipped with one of these norms, \mathcal{B}_γ is a Banach space.*

Proof. The fact that $(\mathcal{B}_\gamma, N_{\infty,\gamma})$ is a Banach space is well-known.

- $N_{\infty,\gamma}$ and $N_{\infty,\gamma,\tilde{\gamma}}$ are equivalent. Since $|f|_{\tilde{\gamma}} \leq |f|_\gamma$, we have $N_{\infty,\gamma,\tilde{\gamma}}(f) \leq N_{\infty,\gamma}(f)$. Conversely, for $x \in M$,

$$\frac{|f(x)|}{p(x)^{\gamma+1}} \leq \frac{|f(x_0)| + \frac{1}{\lambda_0} m_\gamma(f) p(x)^{\gamma+1}}{p(x)^{\gamma+1}} \leq |f(x_0)| + \frac{1}{\lambda_0} m_\gamma(f).$$

The bounds $|f(x_0)| \leq |f|_{\tilde{\gamma}}$ and $1 \leq \lambda_0^{-1}$ prove that $|f|_\gamma \leq \lambda_0^{-1} N_{\infty,\gamma,\tilde{\gamma}}(f)$, consequently $N_{\infty,\gamma}(f) \leq (1 + \lambda_0^{-1}) N_{\infty,\gamma,\tilde{\gamma}}(f)$.

To establish that $N_\gamma(\cdot)$ and $N_{\infty,\gamma}(\cdot)$ are equivalent, we proceed as in Hennion and Hervé (2001) (Chap. X).

Lemma 5.3. *$(\mathcal{B}_\gamma, N_\gamma)$ is a Banach space.*

Proof. Let $(f_n)_n$ be a Cauchy sequence in $(\mathcal{B}_\gamma, N_\gamma)$. Set $g_n = f_n - f_n(y_0)$, where y_0 is any point of M . We have $|g_q(x) - g_p(x)| \leq m_\gamma(g_q - g_p) \Delta_\gamma(x, y_0) = m_\gamma(f_q - f_p) \Delta_\gamma(x, y_0)$ because $g_n(y_0) = 0$. Hence $\nu(|g_q - g_p|) \leq \nu(\Delta_\gamma(\cdot, y_0)) m_\gamma(f_q - f_p)$. Recall that $\nu(p^{\gamma+1}) < +\infty$, so that $\nu(\Delta_\gamma(\cdot, y_0)) < +\infty$.

Consequently $(g_n)_n$ is a Cauchy sequence in the Lebesgue space $\mathbb{L}^1(\nu)$; therefore it converges in this space, and $(\nu(g_n))_n$ converges in \mathbb{C} . Moreover, $(\nu(f_n))_n$ converges in \mathbb{C} because, by assumption, it is a Cauchy sequence. It follows that $(f_n(y_0))_n$ converges to a complex number, say $f(y_0)$, and then that $(f_n)_n$ converges in $\mathbb{L}^1(\nu)$. Because y_0 is arbitrary, $(f_n)_n$ converges pointwise to f . We have $\lim_{n \rightarrow +\infty} \nu(f - f_n) = 0$. The properties $f \in \mathcal{B}_\gamma$ and $\lim_{n \rightarrow +\infty} m_\gamma(f - f_n) = 0$ are obtained by standard arguments. \square

- N_γ and $N_{\infty,\gamma}$ are equivalent. For $f \in \mathcal{B}_\gamma$, we have $|\nu(f)| \leq \nu(|f|) \leq |f|_\gamma \nu(p^{\gamma+1})$. Thus $N_\gamma(f) \leq (1 + \nu(p^{\gamma+1})) N_{\infty,\gamma}(f)$. Since $(\mathcal{B}_\gamma, N_\gamma)$ and $(\mathcal{B}_\gamma, N_{\infty,\gamma})$ are Banach spaces, the open mapping theorem yields the claimed equivalence, see Dunford and Schwartz (1958).

- $N_{1,\gamma}$ and $N_{\infty,\gamma}$ are equivalent. We have $|f(y)| \leq |f(x)| + m_\gamma(f) d(x, y) p(x)^\gamma p(y)^\gamma$ for all

$x, y \in M$. By integrating this inequality with respect to the measure ν , we obtain

$$\begin{aligned} |f(y)| &\leq \nu(|f|) + m_\gamma(f)p(y)^\gamma \int_M d(x, x_0)p(x)^\gamma d\nu(x) \\ &\quad + m_\gamma(f)d(y, x_0)p(y)^\gamma \int_M p(x)^\gamma d\nu(x) \\ &\leq \nu(|f|) + 2\lambda_0^{-1}m_\gamma(f)p(y)^{\gamma+1}\nu(p^{\gamma+1}), \end{aligned}$$

hence $N_{\infty, \gamma}(f) \leq (1 + 2\lambda_0^{-1}\nu(p^{\gamma+1}))N_{1, \gamma}(f)$.

Finally we have $N_{1, \gamma}(f) = m_\gamma(f) + \nu(|f|) \leq m_\gamma(f) + |f|_\gamma \nu(p^{\gamma+1}) \leq (1 + \nu(p^{\gamma+1}))N_{\infty, \gamma}(f)$. \square

We conclude this subsection by giving a statement that will be useful for the spectral study of $P(t)$.

Lemma 5.4.

- (i) For $0 < \gamma < \tilde{\gamma}$, the canonical embedding from $(\mathcal{B}_\gamma, N_{\infty, \gamma, \tilde{\gamma}})$ into $(\mathcal{B}_\gamma, |\cdot|_{\tilde{\gamma}})$ is compact.
- (ii) For $\gamma \in]0, \gamma_0]$, the canonical embedding from $(\mathcal{B}_\gamma, N_{1, \gamma})$ into $(\mathcal{B}_\gamma, \nu(|\cdot|))$ is compact.

Proof. (i) Let $(f_n)_n$ be a sequence of functions in \mathcal{B}_γ such that $N_{\infty, \gamma, \tilde{\gamma}}(f_n) \leq 1$ for all n . Then $(f_n)_n$ is equicontinuous on every compact set of M , and the diagonal process ensures that there exists a subsequence $(f_{\phi(n)})_n$ which converges uniformly on every compact set of M to a function $f \in \mathcal{B}_\gamma$ satisfying $N_{\infty, \gamma, \tilde{\gamma}}(f) \leq 1$. To prove (i), it suffices now to show that $\lim_n |f - f_{\phi(n)}|_{\tilde{\gamma}} = 0$. Observe that $|f - f_n|_\gamma \leq \lambda_0^{-1}N_{\infty, \gamma, \tilde{\gamma}}(f - f_n) \leq 2\lambda_0^{-1}$ (proof of Prop. 5.2). Let $\varepsilon > 0$. As $\gamma < \tilde{\gamma}$, there exists a positive constant c such that, for all $n \in \mathbb{N}$ and for all $x \in M$ satisfying $d(x, x_0) > c$, we have $\frac{|f(x) - f_n(x)|}{p(x)^{\tilde{\gamma}+1}} \leq \frac{2\lambda_0^{-1}p(x)^{\gamma+1}}{p(x)^{\tilde{\gamma}+1}} \leq \varepsilon$. Besides, on the compact set $M_c = \{x : x \in M, d(x, x_0) \leq c\}$, $(f_{\phi(n)})_n$ converges uniformly to f , thus there exists $N \in \mathbb{N}$ such that, for all $n \geq N$ and all $x \in M_c$, we have $\frac{|f(x) - f_{\phi(n)}(x)|}{p(x)^{\tilde{\gamma}+1}} \leq \varepsilon$. Consequently, for $n \geq N$, we obtain $|f - f_{\phi(n)}|_{\tilde{\gamma}} \leq \varepsilon$.

(ii) Now let $(f_n)_n$ be a sequence of functions in \mathcal{B}_γ such that $N_{1, \gamma}(f_n) \leq 1$. Since $N_{1, \gamma}$ and $N_{\infty, \gamma}$ are equivalent (Prop. 5.2), the sequence $(f_n)_n$ is bounded in $(\mathcal{B}_\gamma, N_{\infty, \gamma})$ by a constant c' . As above we can check that there exists a subsequence $(f_{\phi(n)})_n$ which converges pointwise to a function $f \in \mathcal{B}_\gamma$. Since $|f_n| \leq c'p^{\gamma+1}$ and $p^{\gamma+1}$ is ν -integrable, the Lebesgue theorem ensures that $\lim_n \nu(|f - f_{\phi(n)}|) = 0$. \square

5.3 Quasi-compactness of P on \mathcal{B}_γ

The following statement shows that, for $\gamma \in]0, \gamma_0]$, P is a quasi-compact operator on \mathcal{B}_γ . This property will also follow from arguments given in Section 7 ; but Theorem 5.5 below provides a precise description of the peripheral spectrum of P : 1 is a simple eigenvalue and it is the unique peripheral spectral value of P .

Theorem 5.5. For every $\gamma \in]0, \gamma_0]$, P is a bounded linear operator on \mathcal{B}_γ , and we have the following decomposition

$$\mathcal{B}_\gamma = (\mathbf{C} \cdot 1) \oplus H_\gamma,$$

where $H_\gamma = \{f : f \in \mathcal{B}_\gamma, \nu(f) = 0\}$ is a closed P -invariant subspace of \mathcal{B}_γ such that $r(P|_{H_\gamma}) \leq (\vartheta_0)^{\frac{1}{n_0}} < 1$; the real number $\vartheta_0 < 1$ has been defined in Lemma 5.1, and $r(P|_{H_\gamma})$ is the spectral radius of the restriction of P to H_γ .

Proof. Here it is convenient to consider \mathcal{B}_γ equipped with the norm N_γ . We have for all $k \geq 1$

$$\begin{aligned} \int_G \frac{\Delta_\gamma(gx, gy)}{\Delta_\gamma(x, y)} d\pi^{*k}(g) &= \int_G \frac{d(gx, gy)}{d(x, y)} \left(\frac{p(gx)}{p(x)} \right)^\gamma \left(\frac{p(gy)}{p(y)} \right)^\gamma d\pi^{*k}(g) \\ &\leq \int_G c(g) \delta(g)^{2\gamma} d\pi^{*k}(g) = D_k(\gamma). \end{aligned}$$

Since $\delta \leq \tilde{\delta}$, and c and $\tilde{\delta}$ are submultiplicative (Lemma 4.1), Hypothesis $\mathcal{M}'_{2\gamma_0+1} < +\infty$ and Fubini's theorem ensure that $D_k(\gamma) < +\infty$. Let $f \in \mathcal{B}_\gamma$. We have for $x, y \in M$,

$$\begin{aligned} |P^k f(x) - P^k f(y)| &\leq \int_G |f(gx) - f(gy)| d\pi^{*k}(g) \\ &\leq m_\gamma(f) \Delta_\gamma(x, y) \int_G \frac{\Delta_\gamma(gx, gy)}{\Delta_\gamma(x, y)} d\pi^{*k}(g) \leq m_\gamma(f) \Delta_\gamma(x, y) D_k(\gamma). \end{aligned}$$

With $k = 1$ the previous proves that $Pf \in \mathcal{B}_\gamma$, and $m_\gamma(Pf) \leq D_1(\gamma) m_\gamma(f)$. Since $\nu(Pf) = \nu(f)$, we see that P is a bounded linear operator on $(\mathcal{B}_\gamma, N_\gamma)$. As $\nu(p^{\gamma+1}) < +\infty$, the distribution ν defines a continuous linear functional on \mathcal{B}_γ , consequently $H_\gamma = \text{Ker } \nu$ is a closed subspace; it is P -invariant because $\nu P = \nu$.

On the other hand, with $k = n_0$, since $D_{n_0}(\gamma) \leq \vartheta_0$ (Lemma 5.1), we get $m_\gamma(P^{n_0} f) \leq \vartheta_0 m_\gamma(f)$, and by induction $m_\gamma(P^{qn_0} f) \leq \vartheta_0^q m_\gamma(f)$ for every $q \geq 0$. In particular, if $h \in H_\gamma$, then, for every $q \geq 1$, we have $\nu(P^{qn_0} h) = \nu(h) = 0$, thus $N_\gamma(P^{qn_0} h) = m_\gamma(P^{qn_0} h) \leq \vartheta_0^q m_\gamma(h) = \vartheta_0^q N_\gamma(h)$. Thus $r(P|_{H_\gamma}) = (r(P|_{H_\gamma}^{n_0}))^{\frac{1}{n_0}} \leq (\vartheta_0)^{\frac{1}{n_0}}$.

The identity $f = \nu(f) \cdot 1 + (f - \nu(f) \cdot 1)$ leads to the stated decomposition. \square

6 Fourier operators on \mathcal{B}_γ

Recall that the Fourier kernels $P(t)$, $t \in \mathbb{R}$, associated to P and ξ are defined by

$$(P(t)f)(x) = \int_G e^{it\xi(g,x)} f(gx) d\pi(g),$$

and that ξ is a real valued function on $G \times M$ satisfying Condition (RS) of Section 2.

We shall prove that, for suitable η' , $P(t)$ acts continuously on $\mathcal{B}_{\eta'}$. But, for $0 < \eta' < \eta$, it will also be convenient to see $P(t)$ as a bounded linear map from $\mathcal{B}_{\eta'}$ to \mathcal{B}_{η} ; this is true by virtue of the following topological embedding that will be exploited repeatedly in the sequel :

$$\text{if } 0 < \eta' < \eta, \text{ we have } \mathcal{B}_{\eta'} \subset \mathcal{B}_{\eta}, \quad \text{and, for all } f \in \mathcal{B}_{\eta'}, \quad N_{\infty, \eta}(f) \leq N_{\infty, \eta'}(f).$$

Let $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_{\eta})$ be the space of all bounded linear maps from $(\mathcal{B}_{\eta'}, N_{\infty, \eta'})$ to $(\mathcal{B}_{\eta}, N_{\infty, \eta})$. We denote by $\|\cdot\|_{\eta', \eta}$ the operator norm on $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_{\eta})$; when $\eta' = \eta$, we merely set $\|\cdot\|_{\eta} = \|\cdot\|_{\eta, \eta}$.

6.1 Preliminary remarks about the function $P(\cdot)$

As already mentioned in Section 2, the spectral method described in Hennion and Hervé (2001) consists in applying perturbation theory to $P(t)$, so that the map $P(\cdot)$ has to be sufficiently regular.

In order to understand what are the restrictions imposed here by this property, suppose that Condition (RS) holds with $r > 0$, and let us study the quantity $|P(t)f - Pf|_{\gamma}$ for $f \in \mathcal{B}_{\gamma}$. Let $\varepsilon \in]0, 1]$. From the inequality $|e^{iu} - 1| \leq 2|u|^{\varepsilon}$, Condition (RS), and Lemma 4.1, we have for all $x \in M$,

$$\begin{aligned} |P(t)f(x) - Pf(x)| &\leq \int_G |e^{it\xi(g,x)} - 1| |f(gx)| d\pi(g) \\ &\leq 2|t|^{\varepsilon} (1 + d(x, x_0))^{r\varepsilon} |f|_{\gamma} p(x)^{\gamma+1} \int_G R(g)^{\varepsilon} \frac{p(gx)^{\gamma+1}}{p(x)^{\gamma+1}} d\pi(g) \quad (I') \\ &\leq 2C|t|^{\varepsilon} |f|_{\gamma} p(x)^{\gamma+1} (1 + d(x, x_0))^{r\varepsilon} \quad (I) \end{aligned}$$

with $C = \int_G R(g)^{\varepsilon} \delta(g)^{\gamma+1} d\pi(g)$. Because $(1 + d(\cdot, x_0))^{r\varepsilon}$ is not bounded on M , this estimation does not imply that $\lim_{t \rightarrow 0} |P(t)f - Pf|_{\gamma} = 0$. Similar complications appear when one considers $m_{\gamma}(P(t)f - Pf)$.

To get round these difficulties in the special case of autoregressive processes (§ 3), Milhaud and Raugi (1989) have used a space of locally Lipschitz functions similar to \mathcal{B}_{γ} , which is defined by replacing $p(\cdot)^{\gamma+1}$ with $p(\cdot)^{\gamma+1} e^{\lambda d(\cdot, x_0)}$, where λ is a positive parameter. In this case, provided that the strict contraction and exponential moment conditions given in the above mentioned paper are satisfied, one can verify that the right member of (I') is bounded, and more generally, that $P(\cdot)$ is a regular function from a neighbourhood of $t = 0$ to $\mathcal{L}(\mathcal{B}_{\gamma})$.

In this paper, we use another method which enables us to weaken the contraction and moment hypotheses considered in previous papers; this method is based on the two following facts :

1. By integrating (I) with respect to the measure ν , we obtain $\nu(|P(t)f - Pf|) \leq C'|t|^{\varepsilon} N_{1, \gamma}(f)$. This weak continuity property will be sufficient to apply to $P(t)$ a perturbation theorem of Keller and Liverani (1999).

2. Let $0 < \eta' < \eta$. For $f \in \mathcal{B}_{\eta'}$, we have

$$|P(t)f(x) - Pf(x)| \leq 2C\lambda_0^{-r\varepsilon} |t|^\varepsilon |f|_{\eta'} p(x)^{\eta'+r\varepsilon+1},$$

so that, if $\eta' + r\varepsilon \leq \eta$, we get $|P(t)f - Pf|_\eta \leq 2C\lambda_0^{-r\varepsilon} |t|^\varepsilon |f|_{\eta'}$. This leads to investigate the continuity and, more generally, the existence of Taylor's expansions of $P(t)$ at $t = 0$ when $P(\cdot)$ is viewed as a $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ -valued map (instead of a $\mathcal{L}(\mathcal{B}_{\eta'})$ -valued map) ; this is the aim of Sections 6.2 and 6.3. Let us mention that similar methods are used in Le Page (1989) and Hennion (1991) for other purposes.

6.2 Taylor's expansions of $P(t)$ at $t = 0$

For $\tau > 0$ and any non-negative measurable functions U, V on G , we set

$$\mathcal{I}^\tau(U, V) = \int_G U(g) c(g) \delta(g)^{2\tau} d\pi(g) + \int_G V(g) \delta(g)^{\tau+1} d\pi(g).$$

$\mathcal{I}^\tau(U, V)$ is an additive positively homogeneous function of both U and V , and an increasing function of the variable τ because $\delta(\cdot) \geq 1$.

Observe that, for $0 < \gamma \leq \gamma_0$, we have $\mathcal{I}^\gamma(1, 1) \leq \mathcal{M}'_{2\gamma_0+1} + \mathcal{M}_{\gamma_0+1} < +\infty$.

Let us state the three main results of this section.

Proposition 6.1. *Suppose $s + 1 \leq \gamma_0$, and let γ be a real number such that $s + 1 \leq \gamma \leq \gamma_0$ and*

$$\mathcal{I}^\gamma(0, S) = \int_G S(g) \delta(g)^{\gamma+1} d\pi(g) < +\infty.$$

Then, for all $t \in \mathbb{R}$, $P(t) \in \mathcal{L}(\mathcal{B}_\gamma)$. Besides there exists a constant C such that we have, for all $f \in \mathcal{B}_\gamma$,

$$|P(t)^{n_0} f|_\gamma \leq \mathcal{I}^\gamma(0, 1) |f|_\gamma, \quad m_\gamma(P(t)^{n_0} f) \leq \vartheta_0 m_\gamma(f) + C |t| \mathcal{I}^\gamma(0, S) |f|_\gamma,$$

where $\vartheta_0 < 1$ is the real number defined in Lemma 5.1.

Proposition 6.2. *Suppose that the following condition holds*

$$\mathcal{U}_0(\eta', \eta) : \quad 0 < \eta' \leq \gamma_0, \quad \eta' < \eta, \quad s + 1 \leq \eta, \quad \mathcal{I}^{\eta'}(0, S) < +\infty.$$

Then

$$\lim_{|t| \rightarrow 0} \|P(t) - P\|_{\eta', \eta} = 0.$$

With the view of obtaining Taylor's expansions of $P(t)$ at $t = 0$, let us introduce, for $k \in \mathbb{N}^*$, the kernels

$$(L_k f)(x) = \int_G (i\xi(g, x))^k f(gx) d\pi(g).$$

Proposition 6.3. *Let $n \geq 1$. Suppose that the following condition holds*

$$\mathcal{U}_n(\eta', \eta) : 0 < \eta' \leq \gamma_0, \quad \eta' + nr < \eta, \quad s+1 + (n-1)r < \eta, \quad \mathcal{I}'(R^n, (R+S)R^{n-1}) < +\infty.$$

Then, for $k = 1, \dots, n$, $L_k \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ and

$$\lim_{|t| \rightarrow 0} \frac{1}{|t|^n} \left\| P(t) - P - \sum_{k=1}^n \frac{t^k}{k!} L_k \right\|_{\eta', \eta} = 0.$$

6.3 Proofs of Propositions 6.1-6.2-6.3

The main tool is Lemma 6.4 below which will be stated in the next technical context.

Let $k \in \mathbb{N}^*$. Consider a complex valued measurable function q on $G^k \times M$.

Let $\alpha, \beta \in \mathbb{R}_+$, and let A, B be non-negative measurable functions on $G^k \times M$. We shall say that the inequalities **(AB)** are satisfied if, for all $h \in G^k$ and for all $(x, y) \in M^2$ satisfying $d(x_0, y) \leq d(x_0, x)$, we have

$$(A) \quad |q(h, x)| \leq A(h, x) p(x)^\alpha$$

$$(B) \quad |q(h, x) - q(h, y)| \leq B(h, x) d(x, y) p(x)^\beta.$$

For $x \in M$, we denote by A_x and B_x the non-negative functions defined on G^k by $A_x(h) = A(h, x)$ and $B_x(h) = B(h, x)$.

For $h = (h_1, \dots, h_k) \in G^k$, we set $h^* = h_1 \cdots h_k$, and we denote by $\pi^{\otimes k}$ the product measure on G^k . If $x \in M$ and if f is a measurable function on M such that $h \mapsto q(h, x) f(h^* x)$ is $\pi^{\otimes k}$ -integrable, then we set

$$(\mathcal{K}f)(x) = \int_{G^k} q(h, x) f(h^* x) d\pi^{\otimes k}(h).$$

For $\tau > 0$ and for any non-negative measurable functions U, V on G^k , we set

$$\mathcal{I}_k^\tau(U, V) = \int_{G^k} U(h) c(h^*) \delta(h^*)^{2\tau} d\pi^{\otimes k}(h) + \int_{G^k} V(h) \delta(h^*)^{\tau+1} d\pi^{\otimes k}(h).$$

This integral only occurs in the following technical lemma ; notice that it equals $\mathcal{I}^\tau(U, V)$ when $k = 1$.

Lemma 6.4. *Let $0 < \eta' \leq \eta$. Suppose that, for all $x \in M$, we have $\mathcal{I}_k^{\eta'}(A_x, A_x + B_x) < +\infty$. Then, for $f \in \mathcal{B}_{\eta'}$ and $x \in M$, $\mathcal{K}f(x)$ is defined ; moreover, for $x, y \in M$ such that $x \neq y$ and $d(y, x_0) \leq d(x, x_0)$, we have the inequalities*

$$\begin{aligned} \frac{|\mathcal{K}f(x)|}{p(x)^{\eta+1}} &\leq \frac{\mathcal{I}_k^{\eta'}(0, A_x)}{p(x)^{\eta-\eta'-\alpha}} |f|_{\eta'} \\ \frac{|\mathcal{K}f(x) - \mathcal{K}f(y)|}{\Delta_\eta(x, y)} &\leq \frac{\mathcal{I}_k^{\eta'}(A_x, 0)}{p(x)^{\eta-\eta'-\alpha}} m_{\eta'}(f) + \frac{\mathcal{I}_k^{\eta'}(0, B_x)}{p(x)^{\eta-\beta-1}} |f|_{\eta'}. \end{aligned}$$

To apply this lemma, it will be worth noticing that, for $\eta > 0$ and for any function f on M , we have, owing to symmetry, $m_\eta(f) = \sup \left\{ \frac{|f(x) - f(y)|}{\Delta_\eta(x, y)}, x, y \in M, x \neq y, d(y, x_0) \leq d(x, x_0) \right\}$.

Proof of the lemma. We shall use the inequalities $\sup_{x \in M} \frac{p(gx)}{p(x)} \leq \delta(g)$ (Lemma 4.1) and $|f(g \cdot)| \leq |f|_{\eta'} p(g \cdot)^{\eta'+1}$. Let f, x and y as in the statement. We have

$$\begin{aligned} \int_{G^k} |q(h, x) f(h^*x)| d\pi^{\otimes k}(h) &\leq p(x)^\alpha |f|_{\eta'} \int_{G^k} A(h, x) p(h^*x)^{\eta'+1} d\pi^{\otimes k}(h) \\ &\leq p(x)^{\alpha+\eta'+1} |f|_{\eta'} \mathcal{I}_k^{\eta'}(0, A_x). \end{aligned}$$

It follows that $\mathcal{K}f(x)$ is defined and verifies the first stated inequality. To prove the second one, let us write

$$\begin{aligned} |\mathcal{K}f(x) - \mathcal{K}f(y)| &\leq A_1(x, y) + A_2(x, y) \\ \text{with } A_1(x, y) &= \int_{G^k} |q(h, x)| |f(h^*x) - f(h^*y)| d\pi^{\otimes k}(h) \\ A_2(x, y) &= \int_{G^k} |f(h^*y)| |q(h, x) - q(h, y)| d\pi^{\otimes k}(h). \end{aligned}$$

Then

$$\begin{aligned} \frac{A_1(x, y)}{\Delta_\eta(x, y)} &\leq m_{\eta'}(f) p(x)^\alpha \int_{G^k} A(h, x) \frac{d(h^*x, h^*y) p(h^*x)^{\eta'} p(h^*y)^{\eta'}}{d(x, y) p(x)^\eta p(y)^\eta} d\pi^{\otimes k}(h) \\ &\leq m_{\eta'}(f) \left(\frac{p(x)^\alpha}{p(x)^{\eta-\eta'}} \right) \left(\frac{1}{p(y)^{\eta-\eta'}} \right) \int_{G^k} A(h, x) c(h^*) \delta(h^*)^{2\eta'} d\pi^{\otimes k}(h) \\ &\leq \frac{\mathcal{I}_k^{\eta'}(A_x, 0)}{p(x)^{\eta-\eta'-\alpha}} m_{\eta'}(f) \quad (\text{because } p(y)^{\eta-\eta'} \geq 1) \quad (\mathbf{M1}) \end{aligned}$$

Consider now the quantity $A_2(x, y)$. By using the inequality $d(y, x_0) \leq d(x, x_0)$, we obtain

$$\begin{aligned} A_2(x, y) &\leq |f|_{\eta'} d(x, y) p(x)^\beta \int_{G^k} B(h, x) p(h^*y)^{\eta'+1} d\pi^{\otimes k}(h) \\ &\leq |f|_{\eta'} d(x, y) p(x)^\beta p(y)^{\eta'+1} \mathcal{I}_k^{\eta'}(0, B_x) \\ &\leq |f|_{\eta'} d(x, y) p(x)^{\beta+1} p(y)^{\eta'} \mathcal{I}_k^{\eta'}(0, B_x), \quad (\mathbf{M'2}) \end{aligned}$$

and from $p(y)^{\eta'} \leq p(y)^\eta$, we get

$$\frac{A_2(x, y)}{\Delta_\eta(x, y)} \leq \frac{\mathcal{I}_k^{\eta'}(0, B_x)}{p(x)^{\eta-\beta-1}} |f|_{\eta'} \quad (\mathbf{M2}).$$

We conclude by combining (M1) and (M2). □

We shall also need the next bounds.

Lemma 6.5. For $n \in \mathbb{N}$ and $x \in \mathbb{R}$, we set $\phi_n(x) = e^{ix} - \sum_{k=0}^n \frac{(ix)^k}{k!}$.

For all $x, y \in \mathbb{R}$, we have

1. $|\phi_n(x)| \leq 2|x|^n \min\{1, |x|\}$,
2. $|e^{iy} - e^{ix}| \leq |y - x|$,
3. for $n \geq 1$, $|\phi_n(y) - \phi_n(x)| \leq 2|y - x| \left(|x|^{n-1} \min\{1, |x|\} + |y|^{n-1} \min\{1, |y|\} \right)$,

Proof. The assertion 2 is clear, and it implies that $|\phi_0(x)| \leq \min\{2, |x|\} \leq 2 \min\{1, |x|\}$.

Let $n \geq 1$. The Taylor's formula to the orders n and $n - 1$ with integral remainder shows that $|\phi_n(x)| \leq \frac{|x|^{n+1}}{(n+1)!}$ and $|\phi_n(x)| = |\phi_{n-1}(x) - \frac{(ix)^n}{n!}| \leq 2 \frac{|x|^n}{n!}$. Hence

$$|\phi_n(x)| \leq \min\left\{ \frac{2|x|^n}{n!}, \frac{|x|^{n+1}}{(n+1)!} \right\} \leq 2|x|^n \min\{1, |x|\}.$$

Since $\phi'_n(x) = i\phi_{n-1}(x)$ for $n \geq 1$, we have $|\phi_n(x) - \phi_n(y)| \leq |x - y| \sup\{|\phi_{n-1}(t)| : t \in [x, y]\}$. This inequality and point 1 prove Assertion 3. \square

Now let us prove Propositions 6.1-6.3.

Proof of Proposition 6.1. Let $k \in \mathbb{N}^*$. By induction we easily prove that

$$(P(t)^k f)(x) = \int_{G^k} e^{it\xi_k(h,x)} f(h^*x) d\pi^{\otimes k}(h),$$

with $\xi_k(h, x) = \xi(h_k, x) + \xi(h_{k-1}, h_k x) + \dots + \xi(h_1, h_2 \dots h_k x)$, for all $h = (h_1, \dots, h_k) \in G^k$.

Therefore $\mathcal{K} = P(t)^k$ is associated to the kernel $q(h, x) = e^{it\xi_k(h,x)}$. We have $|q(h, x)| = 1$; Condition (RS) and Lemma 4.1 give for $g_1, g_2 \in G$

$$\begin{aligned} |\xi(g_1, g_2x) - \xi(g_1, g_2y)| &\leq S(g_1) d(g_2x, g_2y) (1 + d(g_2x, x_0) + d(g_2y, x_0))^s \\ &\leq \lambda_0^{-s} d(x, y) S(g_1) c(g_2) (p(g_2x) + p(g_2y))^s \\ &\leq \lambda_0^{-s} d(x, y) S(g_1) c(g_2) \delta(g_2)^s (p(x) + p(y))^s, \end{aligned}$$

hence, if $d(x_0, y) \leq d(x_0, x)$, we get

$$|\xi(g_1, g_2x) - \xi(g_1, g_2y)| \leq 2^s \lambda_0^{-s} d(x, y) S(g_1) c(g_2) \delta(g_2)^s p(x)^s.$$

Finally, by using Lemma 6.5 (Ass. 2) and the facts that $\delta(\cdot) \leq \tilde{\delta}(\cdot)$ and that $c(\cdot)$, $\tilde{\delta}(\cdot)$ are submultiplicative (Lemma 4.1), we obtain that $q(h, x)$ verifies the inequalities (AB) with :

$$A(h, x) = 1, \quad \alpha = 0, \quad B(h, x) = 2^s \lambda_0^{-s} |t| B_k(h), \quad \beta = s,$$

where $B_k(h) = \sum_{i=1}^k S(h_i) c(h_{i+1}) \dots c(h_k) \tilde{\delta}(h_{i+1})^s \dots \tilde{\delta}(h_k)^s$. We have

$$\mathcal{I}_k^\eta(A, A + B) = \mathcal{I}_k^\eta(1, 1 + 2^s \lambda_0^{-s} |t| B_k) \leq \mathcal{I}_k^{\eta_0}(1, 1) + 2^s \lambda_0^{-s} |t| \mathcal{I}_k^{\eta_0}(0, B_k).$$

Since $c\delta^{2\gamma_0} \leq c\tilde{\delta}^{2\gamma_0}$, $\delta^{\gamma_0+1} \leq \tilde{\delta}^{\gamma_0+1}$, and the functions $c, \tilde{\delta}$ are submultiplicative, Hypotheses $\mathcal{M}_{\gamma_0+1} < +\infty$, $\mathcal{M}'_{2\gamma_0+1} < +\infty$, and Fubini's theorem imply that $\mathcal{I}_k^{\gamma_0}(1, 1) < +\infty$. Besides we have

$$\mathcal{I}_k^\gamma(0, B_k) \leq \sum_{i=1}^k \int_{G^k} S(h_i) c(h_{i+1}) \cdots c(h_k) \tilde{\delta}(h_{i+1})^s \cdots \tilde{\delta}(h_k)^s \tilde{\delta}(h_1)^{\gamma+1} \cdots \tilde{\delta}(h_k)^{\gamma+1} d\pi^{\otimes k}(h).$$

We have $\tilde{\delta} \leq \frac{2}{\lambda_0} \delta$, and $\int_G S(g) \delta(g)^{\gamma+1} d\pi(g) < +\infty$, thus $\int_G S(g) \tilde{\delta}(g)^{\gamma+1} d\pi(g) < +\infty$. Moreover we have $c(g) \tilde{\delta}(g)^{\gamma+1+s} \leq c(g) \tilde{\delta}(g)^{2\gamma_0}$. It follows from Hypothesis $\mathcal{M}'_{2\gamma_0+1} < +\infty$ and Fubini's theorem that $\mathcal{I}_k^\gamma(0, B_k) < +\infty$.

Now let us apply Lemma 6.4 with $\eta' = \eta = \gamma \leq \gamma_0$.

For $k = 1$, we get for all $f \in \mathcal{B}_\gamma$

$$|P(t)f|_\gamma \leq \mathcal{I}^\gamma(0, 1) |f|_\gamma.$$

On the other hand, since $\gamma \geq s + 1$, we have $p(x)^{\gamma-s-1} \geq 1$, hence, since $B_1 = S$,

$$m_\gamma(P(t)) \leq \mathcal{I}^\gamma(1, 0) m_\gamma(f) + 2^s \lambda_0^{-s} |t| \mathcal{I}^\gamma(0, S) |f|_\gamma.$$

This proves that $P(t) \in \mathcal{L}(\mathcal{B}_\gamma)$.

For $k = n_0$, the first inequality is still valid for $P(t)^{n_0}$, while the second one becomes

$$m_\gamma(P(t)^{n_0} f) \leq \mathcal{I}_{n_0}^\gamma(1, 0) m_\gamma(f) + 2^s \lambda_0^{-s} |t| \mathcal{I}_{n_0}^\gamma(0, B_{n_0}) |f|_\gamma,$$

with $\mathcal{I}_{n_0}^\gamma(1, 0) \leq \mathcal{I}_{n_0}^{\gamma_0}(1, 0) = \int_G c(h) \delta(h)^{2\gamma_0} d\pi^{*n_0}(h) = \vartheta_0$. □

To establish Propositions 6.2 and 6.3, we shall employ the notation

$$\tau(t, g, x) = \min\{1, |t|R(g)(1 + d(x_0, x))^r\}.$$

Lemma 6.6. *Let $\eta > 0$ and let U, V be non-negative measurable functions on G such that $\mathcal{I}^\eta(U, V) < +\infty$. Then, for all $\epsilon > 0$,*

$$\lim_{|t| \rightarrow 0} \left(\sup_{x \in M} \frac{\mathcal{I}^\eta \left(U(\cdot) \tau(t, \cdot, x), V(\cdot) \tau(t, \cdot, x) \right)}{(1 + d(x, x_0))^\epsilon} \right) = 0.$$

Proof. Let $\rho > 0$. We have $\tau \leq 1$ and, for $1 + d(x_0, x) \leq \rho$, we can write $\tau(t, g, x) \leq \min\{1, |t|R(g)\rho^r\} = \tau_\rho(t, g)$. Therefore, comparing ρ with $1 + d(x_0, x)$, we obtain, for all $x \in M$,

$$\frac{\mathcal{I}^\eta \left(U(\cdot) \tau(t, \cdot, x), V(\cdot) \tau(t, \cdot, x) \right)}{(1 + d(x, x_0))^\epsilon} \leq \rho^{-\epsilon} \mathcal{I}^\eta(U, V) + \mathcal{I}^\eta(U(\cdot) \tau_\rho(t, \cdot), V(\cdot) \tau_\rho(t, \cdot)).$$

Since $\lim_{|t| \rightarrow 0} \tau_\rho(t, g) = 0$ and $\tau_\rho \leq 1$, the dominated convergence theorem implies that

$$\limsup_{|t| \rightarrow 0} \left(\sup_{x \in M} \frac{\mathcal{I}^\eta \left(U(\cdot) \tau(t, \cdot, x), V(\cdot) \tau(t, \cdot, x) \right)}{(1 + d(x, x_0))^\epsilon} \right) \leq \rho^{-\epsilon} \mathcal{I}^\eta(U, V).$$

Since ρ is arbitrary, this provides the desired statement. \square

Proof of Proposition 6.2. Let us consider the kernel $q(g, x) = e^{it\xi(g, x)} - 1$, $x \in M, g \in G$, which defines the operator $\mathcal{K}_t = P(t) - P$. By Lemma 6.5 (Ass. 1. with $n = 0$, and then Ass. 2.), we have for $x \in M$ and $g \in G$

$$|q(g, x)| = |e^{it\xi(g, x)} - 1| \leq 2 \min\{1, |t| |\xi(g, x)|\} \leq 2\tau(t, g, x),$$

and, if $d(y, x_0) \leq d(x, x_0)$,

$$|q(g, x) - q(g, y)| \leq |t| |\xi(g, x) - \xi(g, y)| \leq 2^s |t| S(g) d(x, y) (1 + d(x_0, x))^s.$$

Lemma 6.4 applied with $k = 1$, and

$$A(g, x) = 2\tau(t, g, x), \quad \alpha = 0, \quad B(g, x) = 2^s \lambda_0^{-s} |t| S(g), \quad \beta = s,$$

yields

$$\frac{|\mathcal{K}_t f(x)|}{p(x)^{\eta+1}} \leq 2\lambda_0^{\eta'-\eta} \frac{\mathcal{I}^{\eta'}(0, \tau(t, \cdot, x))}{(1 + d(x, x_0))^{\eta-\eta'}} |f|_{\eta'}$$

$$\frac{|\mathcal{K}_t f(x) - \mathcal{K}_t f(y)|}{\Delta_\eta(x, y)} \leq 2\lambda_0^{\eta'-\eta} \frac{\mathcal{I}^{\eta'}(\tau(t, \cdot, x), 0)}{(1 + d(x, x_0))^{\eta-\eta'}} m_{\eta'}(f) + 2^s \lambda_0^{1-\eta} |t| \frac{\mathcal{I}^{\eta'}(0, S(\cdot))}{(1 + d(x, x_0))^{\eta-s-1}} |f|_{\eta'}.$$

Since $\eta - \eta' > 0$, $\eta - s - 1 > 0$, $\mathcal{I}^{\eta'}(0, 1) \leq \mathcal{M}_{\gamma_0+1} < +\infty$, and $\mathcal{I}^{\eta'}(1, 0) \leq \mathcal{M}'_{2\gamma_0+1} < +\infty$, we conclude by using Lemma 6.6 with $(U, V) = (0, 1)$ and $(U, V) = (1, 0)$. \square

Proof of Proposition 6.3. Let us consider the kernel $q(g, x) = e^{it\xi(g, x)} - \sum_{k=0}^n \frac{(it\xi(g, x))^k}{k!}$,

and set $\mathcal{K}_t = P(t) - P - \sum_{k=1}^n \frac{t^k}{k!} L_k$.

Assertion 1 of Lemma 6.5 implies that we have, for $x \in M$ and $g \in G$

$$|q(g, x)| \leq 2|t|^n |\xi(g, x)|^n \min\{1, |t| |\xi(g, x)|\} \leq 2|t|^n R(g)^n (1 + d(x_0, x))^{nr} \tau(t, g, x),$$

while Assertion 3 shows that, for $d(x_0, y) \leq d(x_0, x)$,

$$\begin{aligned} |q(g, x) - q(g, y)| &\leq 2|t|^n |\xi(g, x) - \xi(g, y)| \\ &\times \left(|\xi(g, x)|^{n-1} \min\{1, |t| |\xi(g, x)|\} + |\xi(g, y)|^{n-1} \min\{1, |t| |\xi(g, y)|\} \right) \\ &\leq 2^{s+2} |t|^n S(g) d(x, y) R(g)^{n-1} \left(1 + d(x_0, x)\right)^{s+(n-1)r} \tau(t, g, x). \end{aligned}$$

Therefore the inequalities (AB) hold with $k = 1$, and

$$A(g, x) = 2\lambda_0^{-nr} |t|^n R^n(g) \tau(t, g, x), \quad \alpha = nr,$$

$$B(g, x) = 2^{s+2} \lambda_0^{-\beta} |t|^n S(g) R^{n-1}(g) \tau(t, g, x), \quad \beta = s + (n-1)r.$$

From Lemma 6.4 with $k = 1$, it follows that

$$\begin{aligned} \frac{|\mathcal{K}_t f(x)|}{p(x)^{\eta+1}} &\leq 2\lambda_0^{\eta'-\eta} |t|^n \frac{\mathcal{I}^{\eta'}(0, R^n(\cdot)\tau(t, \cdot, x))}{(1+d(x, x_0))^{\eta-\eta'-nr}} |f|_{\eta'} \\ \frac{|\mathcal{K}_t f(x) - \mathcal{K}_t f(y)|}{\Delta_\eta(x, y)} &\leq 2\lambda_0^{\eta'-\eta} |t|^n \frac{\mathcal{I}^{\eta'}(R^n(\cdot)\tau(t, \cdot, x), 0)}{(1+d(x, x_0))^{\eta-\eta'-nr}} m_{\eta'}(f) \\ &\quad + 2^{s+2} \lambda_0^{1-\eta} |t|^n \frac{\mathcal{I}^{\eta'}(0, S(\cdot) R^{n-1}(\cdot)\tau(t, \cdot, x))}{(1+d(x, x_0))^{\eta-\beta-1}} |f|_{\eta'}. \end{aligned}$$

Since $\eta - \eta' - nr > 0$, $\eta - \beta - 1 = \eta - s - (n-1)r - 1 > 0$, and $\mathcal{I}^{\eta'}(R^n, (R+S)R^{n-1}) < +\infty$, the previous inequalities imply that $\mathcal{K}_t \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$; then, by using Lemma 6.6, we get

$$\lim_{|t| \rightarrow 0} \frac{1}{|t|^n} \|\mathcal{K}_t\|_{\eta', \eta} = 0.$$

Finally it remains to prove that, for $k = 1, \dots, n$, $L_k \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$. This derives from the following : on the one hand, $P, P(t) \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ (Prop. 6.2), and on the other hand, by the above, we have $P(t) - P - \sum_{k=1}^{n'} \frac{t^k}{k!} L_k \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ for $n' = 0, \dots, n$. \square

To end this section, we give an additional statement which completes Proposition 6.1 and will be helpful in the proof of Proposition 7.4.

Proposition 6.7. *Assume $s+1 < \gamma_0$, and let η and $\tilde{\eta}$ be such that $s+1+(\tilde{\eta}-\eta) \leq \eta < \tilde{\eta} < \gamma_0$ and*

$$\mathcal{I}^{\tilde{\eta}}(0, S) < +\infty.$$

Then there exists a constant C such that we have, for all $t \in \mathbb{R}$ and $f \in \mathcal{B}_\eta$,

$$m_\eta(P(t)^{n_0} f) \leq \vartheta_0 m_\eta(f) + C|t| |f|_{\tilde{\eta}}.$$

Proof. First we establish the following with the notations of Lemma 6.4.

Lemma 6.8. *Suppose that the inequalities (AB) hold. Let $0 < \eta < \tilde{\eta}$.*

If $\alpha = 0$, $\beta + 1 + \tilde{\eta} < 2\eta$, and if, for all $x \in M$, $\mathcal{I}_k^\eta(A_x, 0) + \mathcal{I}_k^{\tilde{\eta}}(0, B_x) < +\infty$, then we have, for all $f \in \mathcal{B}_\eta$,

$$m_\eta(\mathcal{K}f) \leq \mathcal{I}_k^\eta(A_x, 0) m_\eta(f) + \mathcal{I}_k^{\tilde{\eta}}(0, B_x) |f|_{\tilde{\eta}}.$$

Proof of Lemma 6.8. Let us write, as in the proof of Lemma 6.4, $|\mathcal{K}f(x) - \mathcal{K}f(y)| \leq A_1(x, y) + A_2(x, y)$, and let us return to inequalities (M1) and (M'2).

With $\eta' = \eta$ and $\alpha = 0$, (M1) gives $\frac{A_1(x, y)}{\Delta_\eta(x, y)} \leq \mathcal{I}_k^\eta(A_x, 0) m_\eta(f)$.

The inequality (M'2) holds for any $\eta' > 0$, in particular it is satisfied with $\eta' = \tilde{\eta}$. Besides, if $d(y, x_0) \leq d(x, x_0)$, we have $p(y)^{\tilde{\eta}} = p(y)^{\tilde{\eta}-\eta}p(y)^\eta \leq p(x)^{\tilde{\eta}-\eta}p(y)^\eta$. Hence $A_2(x, y) \leq |f|_{\tilde{\eta}} d(x, y)p(x)^{\beta+1+\tilde{\eta}-\eta}p(y)^\eta \mathcal{I}_k^{\tilde{\eta}}(0, B_x)$. Since $\beta + 1 + \tilde{\eta} - \eta < \eta$, we obtain

$$\frac{A_2(x, y)}{\Delta_\eta(x, y)} \leq \mathcal{I}_k^{\tilde{\eta}}(0, B_x) |f|_{\tilde{\eta}}.$$

We conclude by combining the two previous bounds. \square

Let us now prove the proposition. Consider the kernel $q(g, x) = e^{it\xi_{n_0}(h, x)}$, $h \in G^{n_0}$, $x \in M$, defining $P(t)^{n_0}$ (see proof of Prop. 6.1) ; it verifies the inequalities (AB) with $k = n_0$, and α, β, A, B given in the proof of Proposition 6.1. Lemma 6.8 applies to this kernel because $\beta + 1 + \tilde{\eta} = s + 1 + \tilde{\eta} \leq 2\eta$, $\mathcal{I}_{n_0}^\eta(A_x, 0) = \mathcal{I}_{n_0}^{\gamma_0}(1, 0) \leq \vartheta_0$, and $\mathcal{I}_{n_0}^{\tilde{\eta}}(0, B_x) < +\infty$; this last point can be shown by using Hypothesis $\mathcal{I}^{\tilde{\eta}}(0, S) < +\infty$ and a method similar to that employed in the proof of Proposition 6.1. This proves the proposition. \square

7 The spectrum of $P(t)$ acting on \mathcal{B}_γ

We use the standard notations $\sigma(T)$ and $r(T)$ to name the spectrum and the spectral radius of an operator T , see Dunford and Schwartz (1958). We denote by \mathcal{B}'_γ the topological dual space of \mathcal{B}_γ , and by $\langle \cdot, \cdot \rangle$ the canonical bilinear functional on $\mathcal{B}'_\gamma \times \mathcal{B}_\gamma$.

For $\gamma \leq \gamma_0$, the P -invariant probability distribution ν defines an element of \mathcal{B}'_γ , and Theorem 5.5 shows that $P \in \mathcal{L}(\mathcal{B}_\gamma)$, that

$$\sigma(P) \subset \{1\} \cup \{z : z \in \mathbb{C}, |z| \leq \kappa_0\}, \quad \text{with } \kappa_0 = \vartheta_0^{\frac{1}{n_0}} < 1.$$

and that there exists $N_{(\gamma)} \in \mathcal{L}(\mathcal{B}_\gamma)$, with spectral radius $r(N_{(\gamma)}) \leq \kappa_0 < 1$, such that, for $n \geq 1$ and $f \in \mathcal{B}_\gamma$,

$$P^n f = \langle \nu, f \rangle 1 + N_{(\gamma)}^n f.$$

The following statement which is obtained by applying to $P(\cdot)$ a perturbation theorem of Keller and Liverani (1999), asserts firstly that, for small $|t|$, the spectrum of $P(t)$ is close to that of P , secondly that a spectral decomposition of the preceding type is still valid for $P(t)$, and thirdly that the resolvents are uniformly bounded in t for z ranging outside a neighbourhood of the spectrum of P .

We shall use the following notations.

Let κ'_0 and κ''_0 be real numbers such that $0 < \kappa_0 < \kappa'_0 < \kappa''_0 < 1$. Let \mathcal{D}_0 and \mathcal{D}_1 be the open discs of the complex plane defined by

$$\mathcal{D}_0 = \{z : z \in \mathbb{C}, |z| < \kappa'_0\} \quad \mathcal{D}_1 = \{z : z \in \mathbb{C}, |z - 1| < 1 - \kappa''_0\}.$$

We denote by Γ_0 and Γ_1 the oriented circles defined respectively as the boundaries of \mathcal{D}_0 and of \mathcal{D}_1^c . We set

$$\mathcal{R} = \mathbb{C} \setminus (\mathcal{D}_0 \cup \mathcal{D}_1) = \{z : z \in \mathbb{C}, |z| \geq \kappa'_0, |z - 1| \geq 1 - \kappa''_0\}.$$

Proposition 7.1. *Assume that $s + 1 \leq \gamma_0$. Let γ be such that $s + 1 \leq \gamma \leq \gamma_0$ and*

$$\mathcal{I}^\gamma(0, S) = \int_G S(g) \delta(g)^{\gamma+1} d\pi(g) < +\infty.$$

Then, for all $t \in \mathbb{R}$, $P(t) \in \mathcal{L}(\mathcal{B}_\gamma)$. Moreover there exists an open interval I_γ containing $t = 0$ such that we have the following spectral properties, for $t \in I_\gamma$, and for $P(t)$ acting on \mathcal{B}_γ :

(a) $\sigma(P(t)) \subset \mathcal{D}_0 \cup \mathcal{D}_1$, and there exists $\lambda_{(\gamma)}(t) \in \mathbb{C}$ such that $\sigma(P(t)) \cap \mathcal{D}_1 = \{\lambda_{(\gamma)}(t)\}$,

(b) there exists a unique function $v_{(\gamma)}(t)$, belonging to \mathcal{B}_γ , such that we have $\langle \nu, v_{(\gamma)}(t) \rangle = 1$ and $P(t)v_{(\gamma)}(t) = \lambda_{(\gamma)}(t)v_{(\gamma)}(t)$,

(c) we have $M_\gamma = \sup \left\{ \|(z - P(t))^{-1}\|_\gamma, t \in I_\gamma, z \in \mathcal{R} \right\} < +\infty$,

(d) there exist $\phi_{(\gamma)}(t) \in \mathcal{B}'_\gamma$ and $N_{(\gamma)}(t) \in \mathcal{L}(\mathcal{B}_\gamma)$ such that

$$\forall f \in \mathcal{B}_\gamma, \quad \forall n \in \mathbb{N}^*, \quad P(t)^n f = \lambda_{(\gamma)}(t)^n \langle \phi_{(\gamma)}(t), f \rangle v_{(\gamma)}(t) + N_{(\gamma)}(t)^n f.$$

with $\|N_{(\gamma)}(t)^n\|_\gamma \leq \frac{M_\gamma}{2\pi} (\kappa'_0)^n$.

Notice that, for $t = 0$, we have $\lambda_{(\gamma)}(0) = 1$, $v_{(\gamma)}(0) = 1$, $\phi_{(\gamma)}(0) = \nu$, and $N_{(\gamma)}(0) = N_{(\gamma)}$. From the inclusion $\mathcal{B}_{\gamma'} \subset \mathcal{B}_\gamma$, for $0 < \gamma' < \gamma$, and from Proposition 7.1, we deduce the following corollary.

Corollary 7.2. *Under the conditions of Proposition 7.1, if $s + 1 \leq \gamma' < \gamma \leq \gamma_0$, then, for all $t \in I_{\gamma'} \cap I_\gamma$, we have*

$$\lambda_{(\gamma)}(t) = \lambda_{(\gamma')}(t), \quad v_{(\gamma)}(t) = v_{(\gamma')}(t), \quad \phi_{(\gamma)}(t)|_{\mathcal{B}_{\gamma'}} = \phi_{(\gamma')}(t), \quad N_{(\gamma)}(t)|_{\mathcal{B}_{\gamma'}} = N_{(\gamma')}(t).$$

Notations. In accordance with this corollary, when Proposition 7.1 applies to $P(t)$ acting on \mathcal{B}_γ , we set

$$\lambda(t) = \lambda_{(\gamma)}(t), \quad v(t) = v_{(\gamma)}(t), \quad \phi(t) = \phi_{(\gamma)}(t), \quad N(t) = N_{(\gamma)}(t).$$

It will follow from the proof of Proposition 7.1 that we have the following :

Corollary 7.2'. *Under the conditions of Proposition 7.1, for $s + 1 \leq \gamma \leq \gamma_0$ and for $t \in I_\gamma$, the elements $N(t)$, $v(t)$, $\phi(t)$ are given by the following formulae in which integration is considered in the space $\mathcal{L}(\mathcal{B}_\gamma)$*

$$N(t) = \frac{1}{2i\pi} \int_{\Gamma_0} (z - P(t))^{-1} dz, \quad v(t) = \frac{1}{\nu(\Pi(t)1)} \Pi(t)1, \quad \phi(t) = \Pi(t)^* \nu,$$

where
$$\Pi(t) = \frac{1}{2i\pi} \int_{\Gamma_1} (z - P(t))^{-1} dz.$$

Moreover we have
$$N(t)^n = \frac{1}{2i\pi} \int_{\Gamma_0} z^n (z - P(t))^{-1} dz \quad \text{and} \quad \|N(t)^n\|_\gamma \leq \frac{M_\gamma}{2\pi} (\kappa'_0)^n.$$

Proof of Proposition 7.1. The hypotheses are those of Proposition 6.1. Consequently, for all $t \in \mathbb{R}$, $P(t) \in \mathcal{L}(\mathcal{B}_\gamma)$.

To establish the assertions (a) to (d), we shall use the results of Keller and Liverani (1999). Let us specify the context of this paper : the space (here \mathcal{B}_γ) on which the collection of operators (here $P(t)$, $t \in \mathbb{R}$) acts, is endowed with a norm (here $N_{1,\gamma}$, § 5.2) with respect to which the space is complete, and with an auxiliary norm which is dominated by the preceding one. An easy adaption shows that the results of Keller and Liverani (1999) are still valid with an auxiliary semi-norm (here $\nu(|\cdot|)$). The lemma below proves that the required hypotheses are fulfilled.

Lemma 7.3. *Under the hypotheses of the proposition :*

1. for $t \in \mathbb{R}$, $n \in \mathbb{N}^*$ and $f \in \mathcal{B}_\gamma$, we have $\nu(|P(t)^n f|) \leq \nu(|f|)$,
2. there exist $J \in \mathbb{R}_+$ and an open interval I_γ containing $t = 0$ such that, for $t \in I_\gamma$, we have

$$\forall f \in \mathcal{B}_\gamma, \quad N_{1,\gamma}(P(t)^{n_0} f) \leq (\kappa'_0)^{n_0} N_{1,\gamma}(f) + J \nu(|f|),$$

3. for all $t \in I_\gamma$, the essential spectral radius of $P(t)$ is $\leq \kappa'_0$,
4. there exists a positive continuous function φ , vanishing at $t = 0$, such that we have, for all $f \in \mathcal{B}_\gamma$, $\nu(|P(t)f - Pf|) \leq \varphi(t) N_{1,\gamma}(f)$.

We refer to Hennion and Hervé (2001), Chap. XIV, for the notion of essential spectral radius of an operator. The property 4 above means that, in a weak sense, for small $|t|$, $P(t)$ is a perturbation of P .

Proof of the lemma. 1. As P is non-negative, we get $|P(t)^n f| \leq P^n |f|$, hence the inequality of point 1, since ν is P -invariant.

2. From Proposition 6.1, we have, for all $f \in \mathcal{B}_\gamma$,

$$m_\gamma(P(t)^{n_0} f) \leq \kappa_0^{n_0} m_\gamma(f) + C|t| \mathcal{I}^\gamma(0, S) |f|_\gamma.$$

As a consequence of the equivalence of the norms $N_{\infty,\gamma}$ and $N_{1,\gamma}$, we get a constant K' , such that, for all $f \in \mathcal{B}_\gamma$, we have

$$m_\gamma(P(t)^{n_0} f) \leq \kappa_0^{n_0} m_\gamma(f) + K'|t| N_{1,\gamma}(f) = \left(\kappa_0^{n_0} + K'|t| \right) m_\gamma(f) + K'|t| \nu(|f|).$$

so that, for $|t| \leq \frac{\kappa_0^{n_0} - \kappa'_0^{n_0}}{K'}$, we obtain

$$m_\gamma(P(t)^{n_0} f) \leq (\kappa'_0)^{n_0} m_\gamma(f) + (\kappa_0^{n_0} - \kappa'_0^{n_0}) \nu(|f|).$$

Using point 1, we get $N_{1,\gamma}(P(t)^{n_0} f) \leq (\kappa'_0)^{n_0} N_{1,\gamma}(f) + J \nu(|f|)$ with $J = \kappa_0^{n_0} - \kappa'_0^{n_0} + 1$.

3. Recall that the essential spectral radius of an operator is smaller than its spectral radius, consequently, point 3 is clear when $r(P(t)) \leq \kappa'_0$.

Assume that $r(P(t)) > \kappa'_0$. Then, from point 1 and the Doeblin-Fortet inequality established in point 2, and from the fact that the canonical embedding of $(\mathcal{B}_\gamma, N_{1,\gamma})$ into $(\mathcal{B}_\gamma, \nu(|\cdot|))$ is compact (Lemma 5.4), we deduce by means of the Ionescu-Tulcea and Marinescu theorem or more precisely of Corollary 1 in Hennion (1993) that, for $|t| \leq \frac{\kappa_0^{m_0} - \kappa_0^{n_0}}{K'}$, $P(t)$ is quasi-compact, and that its essential spectral radius is $\leq \kappa'_0$.

4. Using the inequality $|f(gx)| \leq |f|_\gamma p(gx)^{\gamma+1} \leq |f|_\gamma \tilde{\delta}(g)^{\gamma+1} p(x)^{\gamma+1}$ (Lemma 4.1), we obtain $\nu(|P(t)f - Pf|) \leq \int_G \int_M |e^{it\xi(g,x)} - 1| |f(gx)| d\pi(g) d\nu(x) \leq |f|_\gamma \varepsilon(t)$, with

$$\varepsilon(t) = \int_G \int_M |e^{it\xi(g,x)} - 1| \tilde{\delta}(g)^{\gamma+1} p(x)^{\gamma+1} d\pi(g) d\nu(x).$$

Since $\nu(p^{\gamma+1}) < +\infty$ and $\pi(\tilde{\delta}^{\gamma+1}) = \mathcal{M}_{\gamma+1} \leq \mathcal{M}_{\gamma_0+1} < +\infty$, it follows From Lebesgue's theorem that ε is a continuous function on \mathbb{R} , which vanishes at $t = 0$. Point 4 is deduced from the above inequality and the equivalence of the norms $N_{\infty,\gamma}$ and $N_{1,\gamma}$. \square

Now the assertions (a) and (c) of Proposition 7.1 follow directly from the results of Keller and Liverani (1999) which moreover assert that

$$\Pi_{(\gamma)}(t) = \frac{1}{2i\pi} \int_{\Gamma_1} (z - P(t))^{-1} dz$$

is a rank 1 bounded projection from \mathcal{B}_γ onto $\text{Ker}(P(t) - \lambda(t))$, and that $\nu(|\Pi_{(\gamma)}(t)1 - \Pi_{(\gamma)}(0)1|) = \nu(|\Pi_{(\gamma)}(t)1 - 1|)$ converges to 0 with t .

Therefore, for sufficiently small $|t|$, we have $\nu(\Pi_{(\gamma)}(t)1) \neq 0$, and we can set

$$v_{(\gamma)}(t) = \frac{1}{\nu(\Pi_{(\gamma)}(t)1)} \Pi_{(\gamma)}(t)1,$$

this function verifies the condition (b) of Proposition 7.1. The assertion (d) and Corollary 7.2' also follow from Keller and Liverani (1999). \square

We conclude this section with a result that will be useful for the study of the non-arithmeticity of ξ (cf. § 9).

Proposition 7.4. *Assume that the conditions of Proposition 7.1 are satisfied and reinforced by $s + 1 < \gamma < \gamma_0$ and by the existence of $\tilde{\gamma}$, $\gamma < \tilde{\gamma} < \gamma_0$, such that $\mathcal{I}^{\tilde{\gamma}}(0, S) < +\infty$. Let $t \in \mathbb{R}$ be such that, for $P(t)$ acting on \mathcal{B}_γ , we have $r(P(t)) \geq 1$. Then $r(P(t)) = 1$ and $P(t)$ is quasi-compact on \mathcal{B}_γ .*

Proof. Since $s + 1 < \gamma < \gamma_0$, we can suppose that $\tilde{\gamma}$ verifies $s + 1 + (\tilde{\gamma} - \gamma) \leq \gamma$. For convenience, we set $\tilde{N}(f) = N_{\infty,\gamma,\tilde{\gamma}}(f) = m_\gamma(f) + |f|_{\tilde{\gamma}}$ (§ 5.2).

The first inequality of Proposition 6.1, when applied to $\tilde{\gamma}$ and to the kernel $q(g, x) = e^{it\xi(g,x)}$, shows that

$$|P(t)f|_{\tilde{\gamma}} \leq \mathcal{I}^{\tilde{\gamma}}(0, 1)|f|_{\tilde{\gamma}},$$

with $\mathcal{I}^{\tilde{\eta}}(0, 1) \leq \mathcal{I}^{\eta}(0, 1) < +\infty$. Moreover Proposition 6.7 applied to the couple $(\gamma, \tilde{\gamma}) = (\eta, \tilde{\eta})$ asserts that there exists a constant C such that, for $t \in \mathbb{R}$ and $f \in \mathcal{B}_{\gamma}$, we have

$$m_{\gamma}(P(t)^{n_0} f) \leq \kappa_0^{n_0} m_{\gamma}(f) + C|t| \|f\|_{\tilde{\gamma}}.$$

Setting $C' = C|t| + \mathcal{I}^{\tilde{\eta}}(0, 1)$, we get

$$\tilde{N}(P(t)^{n_0} f) \leq \kappa_0^{n_0} \tilde{N}(f) + C' \|f\|_{\tilde{\gamma}}.$$

From the fact that $P(t)$ is bounded on $(\mathcal{B}_{\gamma}, |\cdot|_{\tilde{\gamma}})$, and since the canonical imbedding of $(\mathcal{B}_{\gamma}, \tilde{N})$ in $(\mathcal{B}_{\gamma}, |\cdot|_{\tilde{\gamma}})$ is compact (Lemma 5.4), we deduce by means of Corollary 1 of Hennion (1993) that, under the condition $r(P(t)) \geq 1$, $P(t)$ is quasi-compact on \mathcal{B}_{γ} , and that its essential spectral radius is $\leq \kappa_0 < \kappa_0'$. Consequently there exists an eigenvalue λ of $P(t)$ such that $|\lambda| = r(P(t))$. Let $w \in \mathcal{B}_{\gamma}$ be an eigenfunction associated with λ . For $n \geq 1$ we have $|\lambda^n w| = |P(t)^n w| \leq P^n |w|$, hence $|\lambda^n| \|w\|_{\gamma} \leq \|P^n |w|\|_{\gamma} \leq \|P^n |w|\|_{\infty, \gamma}$. The spectral decomposition in Theorem 5.5 together with the equivalence of the considered norms on \mathcal{B}_{γ} yield $\sup_n \|P^n |w|\|_{\infty, \gamma} < +\infty$. Hence $|\lambda| \leq 1$, and at last $r(P(t)) = 1$. \square

8 Taylor's expansions for $v(\cdot)$, $\phi(\cdot)$, $N(\cdot)$

The hypotheses in the subsequent statements will imply those of Proposition 7.1 and of its corollaries ; thus, for small $|t|$, the eigenelements of the spectral decomposition described in Proposition 7.1 are defined. We are going to use the Taylor's expansions of $P(\cdot)$ written in Proposition 6.3 to obtain Taylor's expansions for $v(\cdot)$, $\phi(\cdot)$, and $N(\cdot)$.

Proposition 8.1. *(First order Taylor's expansions)*

Suppose that, for $\eta' < \eta$, the following condition holds

$$\mathcal{V}_1(\eta', \eta) : \quad s + 1 \leq \eta' \leq \eta' + r < \eta \leq \gamma_0 \quad \mathcal{I}^{\eta-r}(R, R + S) < +\infty.$$

Then Proposition 7.1 applies to $P(t)$ acting on $\mathcal{B}_{\eta'}$, and the functions $v(\cdot)$, $\phi(\cdot)$, and $N(\cdot)$ from $I_{\eta'}$ in $(\mathcal{B}_{\eta'}, N_{\infty, \eta})$, $\mathcal{B}'_{\eta'}$, and $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_{\eta})$ respectively, have a derivative at $t = 0$. Moreover there exists a constant K_1 such that

$$\forall n \geq 1, \forall t \in I_{\eta'}, \quad \|N(t)^n - N(0)^n\|_{\eta', \eta} \leq K_1 |t| (\kappa_0')^n.$$

Proposition 8.2. *(Second order Taylor's expansions)*

Suppose that, for $\eta' < \eta$, the following condition holds

$$\mathcal{V}_2(\eta', \eta) : \quad s + 1 \leq \eta' \leq \eta' + 2r < \eta \leq \gamma_0, \quad \mathcal{I}^{\eta-r}(R, R + S) + \mathcal{I}^{\eta'}(R^2, (R + S)R) < +\infty.$$

Then Proposition 7.1 applies to $P(t)$ acting on $\mathcal{B}_{\eta'}$, and the functions $v(\cdot)$, $\phi(\cdot)$ and $N(\cdot)$, from $I_{\eta'}$ in $(\mathcal{B}_{\eta'}, N_{\infty, \eta})$, $\mathcal{B}'_{\eta'}$, and $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_{\eta})$ respectively, have second order Taylor's expan-

sions at $t = 0$. Moreover we have, for all $t \in I_{\eta'}$ and $n \geq 1$,

$$N(t)^n = N(0)^n + t N_{1,n} + \frac{t^2}{2} N_{2,n} + t^2 \varepsilon_n(t),$$

with $N_{1,n}, N_{2,n}, \varepsilon_n(t) \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_{\eta})$, $\limsup_{t \rightarrow 0} \sup_{n \geq 1} \|\varepsilon_n(t)\|_{\eta', \eta} = 0$, and $\sup_{n \geq 1} \|N_{j,n}\|_{\eta', \eta} < +\infty$ for $j = 1, 2$.

The rest of this section is devoted to the proofs of these propositions. Recall that $\mathcal{R} = \{z : z \in \mathbb{C}, |z| \geq \kappa'_0, |z - 1| \geq 1 - \kappa''_0\}$. For $\gamma \in]0, \gamma_0]$, we set

$$J_{\gamma} = \sup_{z \in \mathcal{R}} \|(z - P)^{-1}\|_{\gamma} < +\infty \quad (\text{Theorem 5.5}).$$

Under conditions $\mathcal{V}_1(\eta', \eta)$ or $\mathcal{V}_2(\eta', \eta)$, we have $\mathcal{I}^{\eta'}(0, S) \leq \mathcal{I}^{\eta-r}(0, S) < +\infty$ and $s + 1 \leq \eta' \leq \gamma_0$. Consequently Proposition 7.1 applies to $P(t)$ acting on $\mathcal{B}_{\eta'}$. In particular, for $t \in I_{\eta'}$ and for $z \in \mathcal{R}$, $(z - P(t))$ is invertible on $\mathcal{B}_{\eta'}$, and we have

$$M_{\eta'} = \sup \left\{ \|(z - P(t))^{-1}\|_{\eta'}, t \in I_{\eta'}, z \in \mathcal{R} \right\} < +\infty.$$

We shall need the following formula. Let \mathcal{B} be a Banach space. If U and V are bounded operators on \mathcal{B} such that U and $U - V$ are invertible, we have

$$(*) \quad (U - V)^{-1} = \sum_{k=0}^n (U^{-1}V)^k U^{-1} + (U^{-1}V)^{n+1} (U - V)^{-1}.$$

Actually, if $W \in \mathcal{L}(\mathcal{B})$, we have $I - W^{n+1} = \sum_{k=0}^n W^k (I - W)$, and hence, if $I - W$ is invertible,

$$(I - W)^{-1} = \sum_{k=0}^n W^k + W^{n+1} (I - W)^{-1}.$$

The claimed formula follows from the relation $(U - V)^{-1} = (I - U^{-1}V)^{-1} U^{-1}$ and the above equality.

In the proofs below we shall apply (*) with $U = z - P$, $V = P(t) - P$, and thus $U - V = z - P(t)$. Observe that, in the sequel, all the space parameters γ are between $s + 1$ and γ_0 , so that conditions $\mathcal{U}_0(\eta', \eta)$ and $\mathcal{U}_n(\eta', \eta)$, $n \geq 1$, of Propositions 6.2 and 6.3 can be rewritten as

$$\mathcal{U}_0(\eta', \eta) : \eta' < \eta, \mathcal{I}^{\eta'}(0, S) < +\infty.$$

$$\mathcal{U}_n(\eta', \eta) : \eta' + nr < \eta, \mathcal{I}^{\eta'}(R^n, (R + S)R^{n-1}) < +\infty.$$

Otherwise notice that, if $\eta' < \eta_1 < \eta$ and if $T \in \mathcal{L}(\mathcal{B}_{\eta_1}, \mathcal{B}_{\eta})$, then $T \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_{\eta})$ and $\|T\|_{\eta', \eta} \leq \|T\|_{\eta_1, \eta}$.

Proof of Proposition 8.1. The next lemma gives a first order Taylor's expansion for the resolvent $(z - P(t))^{-1}$. We set $R(z, t) = (z - P(t))^{-1}$ and $R(z) = R(z, 0) = (z - P)^{-1}$.

Lemma 8.3. *Under condition $\mathcal{V}_1(\eta', \eta)$, there exists a continuous function R' from \mathcal{R} to $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$, such that we have,*

$$\lim_{t \rightarrow 0} \frac{1}{|t|} \sup_{z \in \mathcal{R}} \left\| (z - P(t))^{-1} - (z - P)^{-1} - tR'_z \right\|_{\eta', \eta} = 0.$$

Proof. Setting n to 1 and U and V to the values indicated a few lines above, the formula (*) gives, for $z \in \mathcal{R}$ and $t \in I_{\eta'}$,

$$R(z, t) = R(z) + R(z)(P(t) - P)R(z) + R(z)(P(t) - P)R(z)(P(t) - P)R(z, t).$$

As, by assumption, $\eta' + r < \eta$, we can choose η_1 such that $\eta' < \eta_1 \leq \eta_1 + r < \eta$. Condition $\mathcal{U}_0(\eta', \eta_1)$ is verified because $\eta' < \eta_1$ and $\mathcal{I}^{\eta'}(0, S) \leq \mathcal{I}^{\eta-r}(0, S) < +\infty$, hence $\lim_{t \rightarrow 0} \|P(t) - P\|_{\eta', \eta_1} = 0$. Condition $\mathcal{U}_1(\eta_1, \eta)$ holds because $\eta_1 + r < \eta$ and $\mathcal{I}^{\eta_1}(R, R + S) \leq \mathcal{I}^{\eta-r}(R, R + S) < +\infty$, hence $P(t) - P = tL_1 + \Upsilon_1(t)$, with $L_1, \Upsilon_1(t) \in \mathcal{L}(\mathcal{B}_{\eta_1}, \mathcal{B}_\eta)$ and $\lim_{t \rightarrow 0} |t|^{-1} \|\Upsilon_1(t)\|_{\eta_1, \eta} = 0$ (Prop. 6.3). Now we write

$$R(z, t) = R(z) + tR'_z + \Theta_1(z, t) + \Theta_2(z, t),$$

with $R'_z = R(z)L_1R(z)$, and

$$\Theta_1(z, t) = R(z)\Upsilon_1(t)R(z), \quad \Theta_2(z, t) = R(z)(P(t) - P)R(z)(P(t) - P)R(z, t).$$

Since $L_1 \in \mathcal{L}(\mathcal{B}_{\eta_1}, \mathcal{B}_\eta) \subset \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ and since $R(\cdot)$ is continuous from \mathcal{R} to both $\mathcal{L}(\mathcal{B}_{\eta'})$ and $\mathcal{L}(\mathcal{B}_\eta)$, R' is continuous from \mathcal{R} to $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$. For $t \in I_{\eta'}$, $0 < |t| \leq 1$, and $z \in \mathcal{R}$, we have

$$|t|^{-1} \|\Theta_1(z, t)\|_{\eta', \eta} \leq |t|^{-1} \|\Theta_1(z, t)\|_{\eta_1, \eta} \leq J_\eta |t|^{-1} \|\Upsilon_1(t)\|_{\eta_1, \eta} J_{\eta_1},$$

$$|t|^{-1} \|\Theta_2(z, t)\|_{\eta', \eta} \leq J_\eta \left(\|L_1\|_{\eta_1, \eta} + |t|^{-1} \|\Upsilon_1(t)\|_{\eta_1, \eta} \right) J_{\eta_1} \|P(t) - P\|_{\eta', \eta_1} M_{\eta'}.$$

The second members do not depend on $z \in \mathcal{R}$ and converge to 0 with t , this proves the lemma. \square

To establish Proposition 8.1, we now use the formulae of Corollary 7.2'. More precisely, the linear maps $\Pi(t)$ and $N(t)$ of the corollary are considered here as elements of $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ since they may be viewed as integrals of functions with values in $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$.

Then Lemma 8.3 shows that $\Pi(\cdot)$ has a derivative at $t = 0$ as a $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ -valued function. Thus $\Pi(\cdot)^*$ has a derivative at $t = 0$ as a $\mathcal{L}(\mathcal{B}'_{\eta'}, \mathcal{B}'_{\eta'})$ -valued function. This proves the first order Taylor's expansions of $v(\cdot)$ and $\phi(\cdot)$. The existence of a derivative for $N(\cdot)$ at $t = 0$ follows in a similar way from Lemma 8.3. On the other hand, from the integral formula $N(t)^n = \frac{1}{2i\pi} \int_{\Gamma_0} z^n (z - P(t))^{-1} dz$, we deduce the existence of a constant K such that, for $n \geq 1$ and $t \in I_{\eta'}$,

$$(\kappa'_0)^{-(n+1)} \|N(t)^n - N(0)^n\|_{\eta', \eta} \leq \sup_{z \in \Gamma_0} \left\| (z - P(t))^{-1} - (z - P)^{-1} \right\|_{\eta', \eta} \leq |t| \left(\sup_{z \in \Gamma_0} \|R'_z\|_{\eta', \eta} + K \right),$$

hence the inequality of Proposition 8.1. \square

Proof of Proposition 8.2. As above, we start with a Taylor's expansion of the resolvent $(z - P(t))^{-1}$.

Lemma 8.4. *Under condition $\mathcal{V}_2(\eta', \eta)$, there exist continuous functions R' et R'' from \mathcal{R} to $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$, such that we have*

$$\lim_{t \rightarrow 0} \frac{1}{t^2} \sup_{z \in \mathcal{R}} \left\| (z - P(t))^{-1} - (z - P)^{-1} - t R'_z - \frac{t^2}{2} R''_z \right\|_{\eta', \eta} = 0.$$

Proof. Retaining the notations of Lemma 8.3 but setting n to 2, the formula (*) gives, for $z \in \mathcal{R}$ and $t \in I_{\eta'}$,

$$\begin{aligned} R(z, t) &= R(z) + R(z)(P(t) - P)R(z) + R(z)(P(t) - P)R(z)(P(t) - P)R(z) \\ &\quad + R(z)(P(t) - P)R(z)(P(t) - P)R(z)(P(t) - P)R(z, t). \end{aligned}$$

Since $\eta' + 2r < \eta$, we can choose η_1 and η_2 such that $\eta' < \eta_1 \leq \eta_1 + r < \eta_2 \leq \eta_2 + r < \eta$. The condition $\mathcal{U}_2(\eta', \eta)$ is verified, hence by Proposition 6.3,

$$P(t) - P = tL_1 + \frac{t^2}{2}L_2 + \Upsilon_2(t),$$

with $P, P(t), L_1, L_2, \Upsilon_2(t) \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$ and $\lim_{t \rightarrow 0} t^{-2} \|\Upsilon_2(t)\|_{\eta', \eta} = 0$.

The conditions $\mathcal{U}_1(\eta_1, \eta_2)$ and $\mathcal{U}_1(\eta_2, \eta)$ are satisfied since we have $\eta_1 + r < \eta_2$, $\eta_2 + r < \eta$ and $\mathcal{I}^n(R, R + S) \leq \mathcal{I}^{\eta_2}(R, R + S) \leq \mathcal{I}^{\eta - r}(R, R + S)$. Then Proposition 6.3 with $n = 1$ shows that

$$P(t) - P = tL_1 + \Upsilon_1(t),$$

with $L_1, \Upsilon_1(t) \in \mathcal{L}(\mathcal{B}_{\eta_1}, \mathcal{B}_{\eta_2}) \cap \mathcal{L}(\mathcal{B}_{\eta_2}, \mathcal{B}_\eta)$ and $\lim_{t \rightarrow 0} |t|^{-1} \|\Upsilon_1(t)\|_{\eta_1, \eta_2} = \lim_{t \rightarrow 0} |t|^{-1} \|\Upsilon_1(t)\|_{\eta_2, \eta} = 0$.

At last, since $\mathcal{I}^{\eta'}(0, S) \leq \mathcal{I}^{\eta - r}(0, S) < +\infty$, the condition $\mathcal{U}_0(\eta', \eta_1)$ holds and Proposition 6.2 ensures that

$$\lim_{t \rightarrow 0} \|P(t) - P\|_{\eta', \eta_1} = 0.$$

We get

$$\begin{aligned} R(z, t) &= R(z) + R(z) \left(tL_1 + \frac{t^2}{2}L_2 + \Upsilon_2(t) \right) R(z) \\ &\quad + R(z) \left(tL_1 + \Upsilon_1(t) \right) R(z) \left(tL_1 + \Upsilon_1(t) \right) R(z) \\ &\quad + R(z) \left(tL_1 + \Upsilon_1(t) \right) R(z) \left(tL_1 + \Upsilon_1(t) \right) R(z) (P(t) - P) R(z, t), \end{aligned}$$

hence

$$R(z, t) = R(z) + tR'_z + \frac{t^2}{2}R''_z + \sum_{k=1}^5 \Theta_k(z, t),$$

with $R'_z = R(z)L_1R(z)$, $R''_z = R(z)L_2R(z) + 2R(z)L_1R(z)L_1R(z)$,

and

$$\Theta_1(z, t) = R(z)\Upsilon_2(t)R(z)$$

$$\Theta_2(z, t) = tR(z)L_1R(z)\Upsilon_1(t)R(z)$$

$$\Theta_3(z, t) = tR(z)\Upsilon_1(t)R(z)L_1R(z)$$

$$\Theta_4(z, t) = R(z)\Upsilon_1(t)R(z)\Upsilon_1(t)R(z)$$

$$\Theta_5(z, t) = R(z)\left(tL_1 + \Upsilon_1(t)\right)R(z)\left(tL_1 + \Upsilon_1(t)\right)R(z)(P(t) - P)R(z, t).$$

Since $L_1 \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta) \cap \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_{\eta_2}) \cap \mathcal{L}(\mathcal{B}_{\eta_2}, \mathcal{B}_\eta)$, $L_2 \in \mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$, and $R(\cdot)$ is continuous from \mathcal{R} to $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$, $\mathcal{L}(\mathcal{B}_{\eta_2})$, and $\mathcal{L}(\mathcal{B}_\eta)$, the functions R' and R'' are continuous from \mathcal{R} to $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$.

We have, for $t \in I$, $0 < |t| \leq 1$, and $z \in \mathcal{R}$

$$t^{-2}\|\Theta_1(z, t)\|_{\eta', \eta} \leq J_\eta(t^{-2}\|\Upsilon_2(t)\|_{\eta', \eta})J_{\eta'},$$

$$t^{-2}\|\Theta_2(z, t)\|_{\eta', \eta} \leq t^{-2}\|\Theta_2(z, t)\|_{\eta_1, \eta} \leq J_\eta\|L_1\|_{\eta_2, \eta}J_{\eta_2}(|t|^{-1}\|\Upsilon_1(t)\|_{\eta_1, \eta_2})J_{\eta_1},$$

$$t^{-2}\|\Theta_3(z, t)\|_{\eta', \eta} \leq t^{-2}\|\Theta_3(z, t)\|_{\eta_1, \eta} \leq J_\eta(|t|^{-1}\|\Upsilon_1(t)\|_{\eta_2, \eta})J_{\eta_2}\|L_1\|_{\eta_1, \eta_2}J_{\eta_1},$$

$$t^{-2}\|\Theta_4(z, t)\|_{\eta', \eta} \leq t^{-2}\|\Theta_4(z, t)\|_{\eta_1, \eta} \leq J_\eta(|t|^{-1}\|\Upsilon_1(t)\|_{\eta_2, \eta})J_{\eta_2}(|t|^{-1}\|\Upsilon_1(t)\|_{\eta_1, \eta_2})J_{\eta_1},$$

$$t^{-2}\|\Theta_5(z, t)\|_{\eta', \eta} \leq J_\eta K_{\eta_2, \eta}J_{\eta_2}K_{\eta_1, \eta_2}J_{\eta_1}\|P(t) - P\|_{\eta', \eta_1}M_{\eta'},$$

with $K_{a,b} = \sup\{\|L_1\|_{a,b} + |t|^{-1}\|\Upsilon_1(t)\|_{a,b}, t \in I, |t| \leq 1\}$.

This proves the lemma because the right hand members do not depend on $z \in \mathcal{R}$ and tend to 0 with t . \square

Let us now complete the proof of Proposition 8.2. The Taylor's expansions for $v(\cdot)$, $\phi(\cdot)$ and $N(\cdot)$ can be deduced from the formulae of Corollaire 7.2'. We just specify how to get the expansion for $N(\cdot)^n$. Using integration in $\mathcal{L}(\mathcal{B}_{\eta'}, \mathcal{B}_\eta)$, we set

$$N_{1,n} = \frac{1}{2i\pi} \int_{\Gamma_0} z^n R'_z dz \quad \text{and} \quad N_{2,n} = \frac{1}{2i\pi} \int_{\Gamma_0} z^n R''_z dz.$$

We have $\|N_{1,n}\|_{\eta', \eta} \leq \kappa_0'^{n+1} \sup_{z \in \Gamma_0} \|R'_z\|_{\eta', \eta}$ and $\|N_{2,n}\|_{\eta', \eta} \leq \kappa_0'^{n+1} \sup_{z \in \Gamma_0} \|R''_z\|_{\eta', \eta}$. Lemma 8.4 yields

$$\begin{aligned} \|\varepsilon_n(t)\|_{\eta', \eta} &= \frac{1}{2\pi t^2} \left\| \int_{\Gamma_0} z^n \left((z - P(t))^{-1} - (z - P)^{-1} - tR'_z - \frac{t^2}{2}R''_z \right) dz \right\|_{\eta', \eta} \\ &\leq \frac{\kappa_0'^{n+1}}{t^2} \sup_{z \in \mathcal{R}} \left\| (z - P(t))^{-1} - (z - P)^{-1} - tR'_z - \frac{t^2}{2}R''_z \right\|_{\eta', \eta}, \end{aligned}$$

since $\kappa_0'^{n+1} \leq 1$, we conclude that $\limsup_{t \rightarrow 0} \sup_{n \geq 1} \|\varepsilon_n(t)\|_{\eta', \eta} = 0$. \square

9 Extensions and proofs of Theorems A-B-C-S

We return to the context of Sections 1 and 2. Theorems A', B', C' below concern the behaviour of the sequence of r.v. $((Z_n, S_n^Z))_n$.

9.1 Theorems A'-B'-C'-S'

Neglecting the technical parameter λ_0 of the preceding sections, we may define \mathcal{B}_γ as the space of locally Lipschitz \mathbf{C} -valued functions f on M such that

$$\ell_\gamma(f) = \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)(1 + d(x, x_0))^\gamma(1 + d(y, x_0))^\gamma}, x, y \in M, x \neq y \right\} < +\infty,$$

endowed with the norm

$$\|f\|_{\infty, \gamma} = \ell_\gamma(f) + \sup_{x \in M} \frac{|f(x)|}{(1 + d(x, x_0))^{\gamma+1}} ;$$

this norm is clearly equivalent to the ones previously defined on \mathcal{B}_γ .

Recall that we set $\tilde{\delta}(g) = 1 + c(g) + d(gx_0, x_0)$. As previously we can omit λ_0 in the definition of the numbers $\mathcal{I}^\tau(U, V)$ (§ 6.2) by replacing now the function δ by $\tilde{\delta}$, that is, by replacing $\mathcal{I}^\tau(U, V)$ by $\mathcal{J}^\tau(U, V)$, already used in Section 2, and defined by

$$\mathcal{J}^\tau(U, V) = \int_G U(g) c(g) \tilde{\delta}(g)^{2\tau} d\pi(g) + \int_G V(g) \tilde{\delta}(g)^{\tau+1} d\pi(g).$$

If $(V, \|\cdot\|)$ is a normed linear space and if $\alpha > 0$, we shall denote by $V(\alpha)$ the closed ball in V with radius α centered at 0. We name $\mathcal{C}_{12}(\mathbb{R})$ the space of \mathbf{C} -valued continuous functions h on \mathbb{R} such that $\lim_{|u| \rightarrow +\infty} u^2 h(u) = 0$.

Under the hypotheses of the next statements, the real number $m = \int_M \int_G \xi(g, x) d\pi(g) d\nu(x)$ is defined, and supposed to be zero.

Recall that Condition $\mathcal{H}(\gamma_0)$ holds if there exist $\gamma_0 \in \mathbb{R}_+^*$ and $n_0 \in \mathbb{N}^*$ such that :

$$\mathcal{M}_{\gamma_0+1} = \pi(\tilde{\delta}^{\gamma_0+1}) < +\infty, \quad \mathcal{M}'_{2\gamma_0+1} = \pi(c \tilde{\delta}^{2\gamma_0}) < +\infty, \quad \mathcal{C}_{2\gamma_0+1}^{(n_0)} = \pi^{*n_0}(c \max\{c, 1\}^{2\gamma_0}) < 1.$$

Theorem A' (Central limit).

Assume $\mathcal{H}(\gamma_0)$ with $\gamma_0 > r + \max\{r, s + 1\}$ and that

$$\int_G R^2 d\pi < +\infty, \quad \mathcal{J}^{\gamma_0-r}(R, R + S) < +\infty.$$

Then, there exists $\sigma^2 \geq 0$ such that, if the r.v. Z satisfies $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$, we have, for $f \geq 0$, $f \in \bigcup_{\gamma < \gamma_0 - r} \mathcal{B}_\gamma$, and for any bounded continuous function h on \mathbb{R} ,

$$\lim_n \mathbb{E} \left[f(Z_n) h \left(\frac{S_n^Z}{\sqrt{n}} \right) \right] = \nu(f) \mathcal{N}(0, \sigma^2)(h).$$

If $h \in C_{\downarrow 2}(\mathbb{R})$, this convergence holds uniformly when (μ, f) ranges over $\mathcal{B}'_{\gamma_0}(\alpha) \times \mathcal{B}_\gamma(\alpha)$.

Theorem B' (Central limit with a rate of convergence).

Assume $\mathcal{H}(\gamma_0)$ with $\gamma_0 > 3r + \max\{r, s + 1\}$ and that

$$\int_G R^3 d\pi < +\infty, \quad \mathcal{J}^{\gamma_0 - r}(R, R + S) + \mathcal{J}^{\gamma_0 - 2r}(R^2, (R + S)R) < +\infty.$$

Then, if $\sigma^2 > 0$, the assertion of Theorem B holds.

Moreover if Z has the distribution ν , then, for $0 < \gamma < \gamma_0 - r$, there exists a positive constant C_γ such that, for $f \in \mathcal{B}_\gamma$, $f \geq 0$, satisfying $\nu(f) > 0$, we have

$$\sup_{u \in \mathbb{R}} \left| \mathbb{E} \left[f(Z_n) 1_{[S_n^Z \leq u\sigma\sqrt{n}]} \right] - \nu(f) \mathcal{N}(0, 1)(] - \infty, u]) \right| \leq \frac{C_\gamma \|f\|_{\infty, \gamma}}{\sqrt{n}}.$$

The statement of the local limit theorem appeals to the non-arithmeticity condition for ξ with respect to the space \mathcal{B}_γ for $\gamma \in]s + 1, \gamma_0 - r[$:

Condition (N-A) $_\gamma$. There is no $t \in \mathbb{R} \setminus \{0\}$, no $\lambda \in \mathbb{C}$, $|\lambda| = 1$, and no bounded function w in \mathcal{B}_γ with non-zero constant modulus on the support Σ_ν of ν , such that we have, for all $x \in \Sigma_\nu$ and for all $n \geq 1$,

$$e^{itS_n^x} w(R_n x) = \lambda^n w(x) \quad \mathbb{P} - a.s.$$

Theorem C' (Local central limit).

Assume that the hypotheses of Theorem A' are satisfied. Let γ be a real number verifying $\max\{r, s + 1\} < \gamma < \gamma_0 - r$ and such that the condition (N-A) $_\gamma$ is fulfilled.

If $\sigma^2 > 0$, and if Z is such that $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$, then for all $f \geq 0$, $f \in \mathcal{B}_\gamma$, and for all $h \in C_{\downarrow 2}(\mathbb{R})$, we have

$$\lim_n \sup_{u \in \mathbb{R}} \left| \sigma \sqrt{2\pi n} \mathbb{E} \left[f(Z_n) h \left(\frac{S_n^Z - u}{\sigma\sqrt{n}} \right) \right] - e^{-\frac{u^2}{2n\sigma^2}} \nu(f) \mathcal{L}(h) \right| = 0,$$

and this convergence holds uniformly when (μ, f) ranges over $\mathcal{B}'_{\gamma_0}(\alpha) \times \mathcal{B}_\gamma(\alpha)$.

Theorem S'.

Assume $\mathcal{H}(\gamma_0)$ with $\gamma_0 > 2r + s + 1$ and that

$$\mathcal{J}^{\gamma_0 - r}(R, R + S) + \mathcal{J}^{\gamma_0 - 2r}(R^2, (R + S)R) < +\infty.$$

Then the assertions of Theorem S hold with $\tilde{\xi}_1 \in \mathcal{B}_{\gamma_0 - r}$ in point (i).

9.2 Proofs of Theorems A'-B'-C'

These proofs are based on expansions of the characteristic function of the r.v S_n^Z .

Proposition 9.1.

(1) Assume that the hypotheses of Theorem A' are fulfilled. Let the parameter γ verifies $\max\{r, s + 1\} < \gamma < \gamma_0 - r$.

Then there exist an open interval I_γ containing $t = 0$, a \mathbf{C} -valued function $\lambda(\cdot)$, and $\mathcal{L}(\mathcal{B}_\gamma)$ -valued functions $L(\cdot)$, $N(\cdot)$, defined on this interval, such that, if the distribution μ of the r.v. Z verifies $\mu(d(\cdot, x_0)^{\gamma_0+1}) < +\infty$, we have, for $n \geq 1$, $t \in I_\gamma$, and for $f \in \mathcal{B}_\gamma$,

$$\mathbb{E}\left[f(Z_n)e^{itS_n^Z}\right] = \langle \mu, P(t)^n f \rangle = \lambda(t)^n \left(\nu(f) + \langle \mu, L(t)f \rangle \right) + \langle \mu, N(t)^n f \rangle.$$

For all $t \in I_\gamma$, we have $|\lambda(t)| \leq 1$, there exists a real positive number $\sigma^2 \geq m^2$ such that

$$\lambda(t) = 1 + imt - \sigma^2 \frac{t^2}{2} + o(t^2),$$

and there exists a positive constant c_γ such that :

(i) if, either $f = 1$ and $\mu \in \mathcal{B}'_{\gamma_0}$, or $f \in \mathcal{B}_\gamma$ and $\mu = \nu$, then

$$|\langle \mu, N(t)^n f \rangle| \leq c_\gamma (\kappa'_0)^n \inf\{|t|, 1\} \|\mu\|_{\infty, \gamma_0} \|f\|_{\infty, \gamma},$$

(ii) $\|N(t)^n\|_\gamma \leq c_\gamma (\kappa'_0)^n$,

(iii) $\|L(t)\|_{\gamma, \gamma_0} \leq c_\gamma \inf\{|t|, 1\}$.

Moreover, if $m = 0$ and $\sigma^2 > 0$, then, for any real number t such that $\frac{t}{\sigma} \in I_\gamma$, we have

(iv) $|\lambda(\frac{t}{\sigma})| \leq e^{-\frac{t^2}{4}}$.

(2) Suppose that the hypotheses of Theorem B' hold. Then, if $m = 0$ and $\sigma^2 > 0$, there exists a constant C_1 such that we have, for all real t such that $\frac{t}{\sigma\sqrt{n}} \in I_\gamma$,

(v) $|\lambda(\frac{t}{\sigma\sqrt{n}})^n - e^{-\frac{t^2}{2}}| \leq \frac{C_1}{\sqrt{n}} |t|^3 e^{-\frac{t^2}{4}}$.

Assume this proposition for a while. To prove Theorems A'- B' -C' we have only to use the method of Hennion and Hervé (2001) Section IV.2 and Chap. VI, which is an adaptation of standard Fourier techniques for sums of i.i.d.r.v. As already mentioned in Section 4.2, we consider here the Fourier kernels $P(t)$ instead of the Fourier kernels $Q(t)$ associated with ξ and the probability transition Q on $G \times M$ defined in Section 4.2. Yet the needed changes are obvious, and we shall not develop the argumentation, we only specify some points.

Firstly, the distribution μ of Z defines an element of \mathcal{B}'_{γ_0} if and only if $\mathbb{E}[d(Z, x_0)^{\gamma_0+1}] < +\infty$, and, in this case, $\|\mu\|_{\infty, \gamma_0} = \mathbb{E}[(1 + d(Z, x_0))^{\gamma_0+1}]$. Actually, we have, for $f \in \mathcal{B}_{\gamma_0}$, $|f| \leq \|f\|_{\infty, \gamma_0} (1 + d(\cdot, x_0))^{\gamma_0+1}$, hence $\mu(|f|) \leq \|f\|_{\infty, \gamma_0} \mathbb{E}[(1 + d(Z, x_0))^{\gamma_0+1}]$.

Secondly, because of the topological embedding of the spaces \mathcal{B}_γ , in the proofs of Theorems A'

and B', it will be sufficient to consider the case where the function f is in a space \mathcal{B}_γ with $\gamma \in]\max\{r, s + 1\}, \gamma_0 - r[$.

At last, in the proof of Theorem C' it is necessary to have some control on the behaviour of $P(t)$, for all $t \in \mathbb{R}$, the following lemma shows how this is related to Condition (N-A) $_\gamma$.

Lemma 9.1.' *Assume conditions of Theorem C' except (N-A) $_\gamma$. Then $P(t)$ is a bounded operator of \mathcal{B}_γ for all $t \in \mathbb{R}$. Let $t \in \mathbb{R}$ such that $r(P(t)) \geq 1$, then there exist $\lambda \in \mathbb{C}$, $|\lambda| = 1$, and a bounded function $w \in \mathcal{B}_\gamma$, with non-zero constant modulus on the support Σ_ν of ν , such that we have, for all $x \in \Sigma_\nu$ and all $n \geq 1$,*

$$e^{itS_n^x}w(R_nx) = \lambda^n w(x) \quad \mathbb{P} - a.s.$$

Consequently, under (N-A) $_\gamma$, for all $t \in \mathbb{R} \setminus \{0\}$, we have $r(P(t)) < 1$.

Proof. Let $\gamma \in]s + 1, \gamma_0 - r[$. The inequality $\mathcal{J}^\gamma(0, S) \leq \mathcal{J}^{\gamma_0 - r}(0, S) < +\infty$ together with Proposition 7.1 shows that the Fourier kernels $P(t)$ act continuously on \mathcal{B}_γ for all $t \in \mathbb{R}$.

By Proposition 7.4, if $r(P(t)) \geq 1$, then $r(P(t)) = 1$, and $P(t)$ is quasi-compact. Consequently, there exist $w \in \mathcal{B}_\gamma \setminus \{0\}$, and $\lambda \in \mathbb{C}$, $|\lambda| = 1$, such that, for all $n \geq 1$, we have

$$P(t)^n w = \lambda^n w.$$

It follows that $|w| \leq P^n |w|$. Since, by Theorem 5.1, the sequence $(P^n |w|)_{n \geq 1}$ converges pointwise to $\nu(|w|)$, we get $|w| \leq \nu(|w|)$, so that w is bounded. From the above and equality $\nu(\nu(|w|)1_M - |w|) = 0$, we deduce that $\nu(\{x : x \in M, |w(x)| = \nu(|w|)\}) = 1$, thus $|w|$ is a non-zero constant function on Σ_ν . For $x \in \Sigma_\nu$ and $n \geq 1$, we write

$$\mathbb{E}\left[1 - \frac{e^{itS_n^x}w(R_nx)}{\lambda^n w(x)}\right] = 1 - \frac{P(t)^n w(x)}{\lambda^n w(x)} = 0.$$

Since $\left|\frac{e^{itS_n^x}w(R_nx)}{\lambda^n w(x)}\right| = 1$, it follows that $e^{itS_n^x}w(R_nx) = \lambda^n w(x) \quad \mathbb{P} - a.s.$ □

To be complete on the properties required for local theorem, one needs to establish the following.

Lemma 9.1''. *Under the conditions of theorem C', for every compact subset K of \mathbb{R}^* ,*

(i) *We have $r_K = \sup\{r(P(t)), t \in K\} < 1$.*

(ii) *There exists $C \geq 0$ and $\rho_K < 1$ such that we have, for all $n \geq 1$, $\sup_{t \in K} \|P(t)^n\| \leq C \rho_K^n$.*

Proof. (i) Suppose that $\sup_{t \in K} r(P(t)) \geq 1$. Then, by Lemma 9.1', $\sup_{t \in K} r(P(t)) = 1$, thus there exists a sequence $(\tau_k)_k$ in K such that $\lim_k r(P(\tau_k)) = 1$. For each $k \geq 1$ consider a spectral value λ_k of $P(\tau_k)$ satisfying $|\lambda_k| = r(P(\tau_k))$. By compactness, one can suppose that $(\tau_k)_k$ and $(\lambda_k)_k$ converge. Set $t_0 = \lim_k \tau_k$, $\lambda = \lim_k \lambda_k$, and observe that $t_0 \in K$, thus $t_0 \neq 0$, and $|\lambda| = 1$.

We are going to show that the perturbation theorem of Keller and Liverani (1999) applies to the action, on a certain space \mathcal{B}_γ , of the family $\{P(t), t \in \mathbb{R}\}$ when $t \rightarrow t_0$. It will follow from this result, see page 145 of the above cited paper, that λ is a spectral value of $P(t_0)$.

But since $t_0 \neq 0$ and $|\lambda| = 1$, this will contradict Lemma 9.1', so we shall get point (i).

Let $\gamma, \tilde{\gamma}$ be such that $s + 1 + (\tilde{\gamma} - \gamma) \leq \gamma < \tilde{\gamma} < \gamma_0 - r$. We establish that $\{P(t), t \in \mathbb{R}\}$ acting on \mathcal{B}_γ satisfies the four assertions of Lemma 7.3, where 0 is replaced by $t_0 \in \mathbb{R}^*$, and the norm $N_{1,\gamma}(\cdot)$ [respectively $\nu(\cdot)$] is replaced by $N_{\infty,\gamma,\tilde{\gamma}}$ [respectively $|\cdot|_{\tilde{\gamma}}$].

1. Using the inequality $|P(t)^n f| \leq P^n |f| \leq |f|_{\tilde{\gamma}} P^n (p^{\tilde{\gamma}+1})$ and Assertion (c) of Lemma 4.2 (observe that $p^{\tilde{\gamma}}$ and $1 + \phi_\lambda$ are equivalent), one easily proves that $\sup_{n \geq 1} |P^n(p^{\tilde{\gamma}+1})|_{\tilde{\gamma}} < +\infty$. It follows that $\{P(t)^n, t \in \mathbb{R}, n \geq 1\}$ is uniformly bounded on $(\mathcal{B}_\gamma, |\cdot|_{\tilde{\gamma}})$.

2. Proposition 6.7 implies the second point of Lemma 7.3 (with $|\cdot|_{\tilde{\gamma}}$ instead of $\nu(|\cdot|)$).

3. If $r(P(t)) > \vartheta_0^{\frac{1}{n_0}}$, where $\vartheta_0 < 1$ is the real number in Proposition 6.7, it follows from Lemma 5.4, from the preceding assertion, and from Hennion (1993), that the essential spectral radius of $P(t)$ is $\leq \vartheta_0^{\frac{1}{n_0}}$. If $r(P(t)) \leq \vartheta_0^{\frac{1}{n_0}}$, this is also valid because the essential spectral radius is always less than the spectral radius.

4. In the same way as Proposition 6.2, it can be proved that there exists a real continuous function $\varepsilon(\cdot)$, vanishing at $t = t_0$, such that we have $\|P(t)f - P(t_0)f\|_{\infty,\tilde{\gamma}} \leq \varepsilon(t) \|f\|_{\infty,\gamma}$ for all $f \in \mathcal{B}_\gamma$. Since $\|\cdot\|_{\infty,\gamma} \leq C \|\cdot\|_{\infty,\gamma,\tilde{\gamma}}$ (Prop. 5.2), we obtain

$$\|P(t)f - P(t_0)f\|_{\tilde{\gamma}} \leq \|P(t)f - P(t_0)f\|_{\infty,\tilde{\gamma}} \leq C \varepsilon(t) \|f\|_{\infty,\gamma,\tilde{\gamma}}.$$

(ii) Let ρ_K be such that $\max\{\vartheta_0^{\frac{1}{n_0}}, r_K\} < \rho_K < 1$, and let Γ be the oriented circle $\{|z| = \rho_K\}$ in \mathbb{C} . For $t \in K$, we have $r(P(t)) \leq r_K < \rho_K$, thus $P(t)^n = \frac{1}{2i\pi} \int_{\Gamma} z^n (z - P(t))^{-1} dz$. Moreover the theorem of Keller-Liverani ensures that, for any $t_0 \in K$, there exists an open interval I , containing t_0 , such that $\sup\{\|(z - P(t))^{-1}\|_{\gamma}, t \in I, |z| = \rho_K\} < +\infty$. By compactness, we get $\sup\{\|(z - P(t))^{-1}\|_{\gamma}, t \in K, |z| = \rho_K\} < +\infty$. This gives (ii). \square

Proof of assertion (1) of Proposition 9.1. Let $\gamma, \max\{r, s + 1\} < \gamma < \gamma_0 - r$.

We have $s + 1 < \gamma < \gamma_0$ and $\mathcal{J}^\gamma(0, S) \leq \mathcal{J}^{\gamma_0-r}(0, S) < +\infty$. Thus Proposition 7.1 applies to $P(t)$ acting on \mathcal{B}_γ . For convenience, the interval I_γ will be denoted by I .

Lemma 9.2. *The maps $v(\cdot)$, $\phi(\cdot)$, and $N(\cdot)$ have derivatives at $t = 0$ as functions with values in $(\mathcal{B}_\gamma, \|\cdot\|_{\infty,\gamma_0})$, \mathcal{B}'_γ , and $\mathcal{L}(\mathcal{B}_\gamma, \mathcal{B}_{\gamma_0})$ respectively, and there exists a constant K such that we have, for all $n \geq 1$ and all $t \in I$,*

$$\|N(t)^n - N(0)^n\|_{\gamma,\gamma_0} \leq K |t| (\kappa'_0)^n.$$

Moreover there exists γ_2 , $0 < \gamma_2 < \gamma$, such that $P(\cdot)$ has a derivative at $t = 0$ as a $\mathcal{L}(\mathcal{B}_{\gamma_2}, \mathcal{B}_\gamma)$ -valued function.

Proof. We have $s + 1 < \gamma \leq \gamma + r < \gamma_0$ and $\mathcal{J}^{\gamma_0-r}(R, R + S) < +\infty$, so that the condition $\mathcal{V}_1(\gamma, \gamma_0)$ is fulfilled and the assertions upon $v(\cdot)$, $\phi(\cdot)$ and $N(\cdot)$ follow from Proposition 8.1. Since $r < \gamma < \gamma_0 - r$, there exists γ_2 such that $0 < \gamma_2 \leq \gamma_2 + r < \gamma \leq \gamma + r < \gamma_0$.

To establish that $P(t)$ has a derivative, we apply Proposition 6.3. Actually, the condition $\mathcal{U}_1(\gamma_2, \gamma)$ holds : we have $\gamma_2 + r < \gamma$, $s + 1 < \gamma$ and $\mathcal{J}^{\gamma_2}(R, R + S) \leq \mathcal{J}^{\gamma_0-r}(R, R + S) < +\infty$. \square

The formula for $\mathbb{E}[e^{itS_n} f(X_n)]$ is obtained by using the basic lemma stated in Section 4.2, the

decomposition of $P(t)$ given in Proposition 7.1, and by setting $L(t)f = \langle \phi(t), f \rangle v(t) - \langle \nu, f \rangle 1$. Under the conditions of (i), we have $\langle \mu, N(0)^n f \rangle = 0$, so that the considered inequality follows from Lemma 9.2.

The inequality (ii) already appears in Corollaire 7.2'.

To obtain (iii), it suffices to remark that, since the functions $v(\cdot)$ and $\phi(\cdot)$ have derivatives in $(\mathcal{B}_\gamma, \|\cdot\|_{\infty, \gamma_0})$ and \mathcal{B}'_γ , there exist constants C_1 and C_2 such that, for $f \in \mathcal{B}_\gamma$,

$$\begin{aligned} \|L(t)f\|_{\infty, \gamma_0} &\leq |\langle \phi(t), f \rangle| \|v(t) - 1\|_{\infty, \gamma_0} + |\langle \phi(t) - \nu, f \rangle| \|1\|_{\infty, \gamma_0} \\ &\leq C_1 |t| \|\phi(t)\|_{\infty, \gamma} \|f\|_{\infty, \gamma} + C_2 |t| \|f\|_{\infty, \gamma} \|1\|_{\infty, \gamma_0}. \end{aligned}$$

It remains to prove the properties of $\lambda(\cdot)$.

From Proposition 7.1, we have $\lambda(0) = 1$ and $\lambda(t)^n = \langle \nu, P(t)^n v(t) \rangle$. Appealing to the invariance of ν , we get $|\lambda(t)|^n \leq \langle \nu, P^n |v(t)| \rangle = \langle \nu, |v(t)| \rangle$. It follows that $|\lambda(t)| \leq 1$.

To prove that $\lambda(\cdot)$ can be expanded to the second order and to identify the terms of its expansion, we proceed as in Lemma IV.4' of Hennion and Hervé (2001).

Lemma 9.3. *For $t \in I$, set $p(t) = \langle \phi(t), 1 \rangle$, $\tilde{v}(t) = \langle \nu, P(t)1 \rangle$, and $u(t) = P(t)1 - \tilde{v}(t)1$. Then $u(0) = 0$, $\langle \nu, u(t) \rangle = 0$, and*

$$\lambda(t) = \frac{1}{p(t)} \langle \phi(t) - \nu, u(t) \rangle + \tilde{v}(t).$$

Proof. The two first equalities are obvious. From the decomposition of Proposition 7.1, we have $P(t)1 = \lambda(t)p(t)v(t) + N(t)1$. As $\langle \phi(t), v(t) \rangle = 1$ and $\phi(t)N(t) = 0$, the formula for $\lambda(t)$ follows from

$$\langle \phi(t), u(t) \rangle = \langle \phi(t), \lambda(t)p(t)v(t) + N(t)1 - \tilde{v}(t)1 \rangle = \lambda(t)p(t) - \tilde{v}(t)p(t). \quad \square$$

Notice that $\tilde{v}(\cdot)$ is the characteristic function of ξ under the distribution $\pi \otimes \nu$, so that the next lemma results from the moment property $\nu(d(\cdot, x_0)^{\gamma_0+1}) < +\infty$.

Lemma 9.4. *Let $n \in \mathbb{N}^*$. Assume that $\int_G R(g)^n d\pi(g) < +\infty$ and that $r \leq \frac{\gamma_0+1}{n}$. Then $\tilde{v}(\cdot)$ has continuous derivatives up to order n , with $\tilde{v}^{(k)}(0) = i^k \int_G \xi(g, x)^k d\pi(g) d\nu(x)$ for $k = 1, \dots, n$.*

We can now obtain the second order Taylor's expansion of $\lambda(\cdot)$.

Lemma 9.5. *$u(\cdot)$ has a derivative at $t = 0$ as a \mathcal{B}_γ -valued function, and we have*

$$\lambda(t) = 1 + imt - \sigma^2 \frac{t^2}{2} + o(t^2), \quad \text{with} \quad \sigma^2 = (\pi \otimes \nu)(\xi^2) - 2\langle \phi'(0), u'(0) \rangle \geq m^2.$$

Proof. By assumption we have $\int_G R(g)^2 d\pi(g) < +\infty$ and $r < \frac{\gamma_0}{2} \leq \frac{\gamma_0+1}{2}$, so that $\tilde{v}(t) = 1 + imt - (\pi \otimes \nu)(\xi^2) \frac{t^2}{2} + o(t^2)$. From Lemma 9.2, we know that $P(\cdot)1$ and $\langle \nu, P(\cdot)1 \rangle$ have derivatives at $t = 0$ as functions with values in \mathcal{B}_γ and \mathbb{C} respectively. Therefore $u(\cdot)$ has a derivative at $t = 0$ as a \mathcal{B}_γ -valued function.

We get, firstly in \mathcal{B}'_γ , $\phi(t) - \nu = \phi(t) - \phi(0) = t\phi'(0) + o(t)$, secondly in \mathcal{B}_γ , $u(t) = tu'(0) + o(t)$, and thirdly in \mathbb{C} , $p(t) = 1 + O(t)$. Setting $c = 2\langle\phi'(0), u'(0)\rangle$, we have

$$\frac{1}{p(t)}\langle\phi(t) - \nu, u(t)\rangle = \left(1 + O(t)\right)\left(c\frac{t^2}{2} + o(t^2)\right) = c\frac{t^2}{2} + o(t^2).$$

We obtain the Taylor's expansion of $\lambda(\cdot)$ by adding the expansion of $\tilde{\nu}$ to the last one. We now prove that $\sigma^2 \geq m$. Setting $\overline{v(t)(\cdot)} = \overline{v(t)(\cdot)}$, we have $P(-t)\overline{v(t)} = \overline{\lambda(t)}\overline{v(t)}$ and, by uniqueness (cf. Prop. 7.1(a)), we get $\lambda(-t) = \overline{\lambda(t)}$. It follows that $\sigma^2 \in \mathbb{R}$. As $1 \geq |\lambda(t)|^2 = 1 - (\sigma^2 - m^2)t^2 + o(t^2)$, we obtain $\sigma^2 - m^2 \geq 0$. Lemma 9.5 is proved. \square

When $m = 0$ and $\sigma^2 > 0$, it follows from the preceding expansion that, for small $|t|$, $|\lambda(\frac{t}{\sigma})| \leq 1 - \frac{t^2}{2} + \frac{t^2}{4} \leq e^{-\frac{t^2}{4}}$, that is (iv).

Proof of the assertion (2) of Proposition 9.1.

The claimed inequality follows (cf. for example Hennion and Hervé (2001)) from the fact that, under the additional hypotheses in (2), the remainder of the second order expansion of $\lambda(\cdot)$ can be specified as follows.

Proposition 9.6. *We have $\lambda(t) = 1 + imt - \sigma^2\frac{t^2}{2} + O(t^3)$.*

Proof.

Lemma 9.7. *There exists $0 < \gamma_2 < \gamma_0$ such that the functions $\phi(\cdot)$ and $P(\cdot)1$ have a second order Taylor's expansion at $t = 0$ as functions with values in \mathcal{B}'_{γ_2} and in \mathcal{B}_{γ_2} respectively.*

Proof. By assumption we have $\gamma_0 > 3r + \max\{r, s + 1\}$, therefore $4r < \gamma_0$ and $s + 1 + r < \gamma_0 - 2r$. It follows that there exist γ_4 and γ_2 such that $0 < \gamma_4 \leq \gamma_4 + 2r < \gamma_2 \leq \gamma_2 + 2r < \gamma_0$ and $s + 1 + r < \gamma_2$.

To establish the assertion on $\phi(\cdot)$, we apply Proposition 8.2, this is possible since the condition $\mathcal{V}_2(\gamma_2, \gamma_0)$ is satisfied, indeed, we have $s + 1 \leq s + 1 + r < \gamma_2 \leq \gamma_2 + 2r < \gamma_0$, and $\mathcal{J}^{\gamma_0-r}(R, R + S) + \mathcal{J}^{\gamma_2}(R^2, (R + S)R) < +\infty$ because $\gamma_2 < \gamma_0 - 2r$.

Moreover, the condition $\mathcal{U}_2(\gamma_4, \gamma_2)$ is verified : we have $0 < \gamma_4 \leq \gamma_4 + 2r < \gamma_2$, $s + 1 + r < \gamma_2$, and then $\mathcal{J}^{\gamma_4}(R^2, (R + S)R) < +\infty$ since $\gamma_4 < \gamma_0 - 2r$. Proposition 6.3 shows that $P(\cdot)$ has a second order Taylor's expansion at $t = 0$ as a $\mathcal{L}(\mathcal{B}_{\gamma_4}, \mathcal{B}_{\gamma_2})$ -valued function, hence the claimed property for $P(\cdot)1$. \square

To conclude, we appeal once more to the formula of Lemma 9.3. Since $\int_G R(g)^3 d\pi(g) < +\infty$ and $r < \frac{\gamma_0}{4} \leq \frac{\gamma_0+1}{3}$, the characteristic function $\tilde{\nu}(\cdot)$ has now three continuous derivatives, so that the remainder of its second order Taylor's expansion is $O(t^3)$. Using the preceding lemma, we have $\phi(t) = \nu + t\phi'(0) + t^2\phi_2 + o(t^2)$ in \mathcal{B}'_{γ_2} and $u(t) = tu'(0) + t^2u_2 + o(t^2)$ in \mathcal{B}_{γ_2} . Consequently $\frac{1}{p(t)}\langle\phi(t) - \nu, u(t)\rangle = (1 + O(t))(c\frac{t^2}{2} + O(t^3)) = c\frac{t^2}{2} + O(t^3)$. It follows that the remainder of the second order expansion of $\lambda(\cdot)$ at $t = 0$ is $O(t^3)$. \square

9.3 Proof of Theorem S'

Proposition 9.8.

Assume $\mathcal{H}(\gamma_0)$ with $\gamma_0 > r + \max\{r, s + 1\}$ and that

$$\int_G R(g)^2 d\pi(g) < +\infty, \quad \mathcal{J}^{\gamma_0-r}(R, R+S) < +\infty,$$

and that $m = 0$.

(a) We set $\theta(x) = \int_G \xi(g, x) d\pi(g)$, $x \in M$. There exists a unique real valued function $w \in \mathcal{B}_{\gamma_0-r}$ such that :

$$\langle \nu, w \rangle = 0 \quad , \quad (1 - P)w = \theta,$$

and we have $\sigma^2 = (\pi \otimes \nu) \left(\xi(\xi + 2w \circ j) \right)$.

(b) Moreover suppose that $\gamma_0 > 2r + s + 1$ and that

$$\mathcal{J}^{\gamma_0-2r}(R^2, (R+S)R) < +\infty.$$

Suppose that the r.v. Z has a distribution μ which defines an element of \mathcal{B}'_{γ_0} . Then, for all $n \geq 1$, the characteristic function $\varphi_n(t) = \mathbb{E}[e^{itS_n^Z}] = \langle \mu, P(t)^{n1} \rangle$ has the Taylor's expansion $\varphi_n(t) = 1 + a_n t + b_n \frac{t^2}{2} + o_n(t^2)$, with $\sup_{n \geq 1} |b_n + n\sigma^2| < +\infty$.

Recall that j defines the action of G on M .

Proof of Proposition 9.8.

The hypothesis $\int_G R(g)^2 d\pi(g) < +\infty$ implies that θ is well defined.

Proof of assertion (a). To begin, we state the differential properties that we shall use.

Let γ be such that $\max\{r, s + 1\} < \gamma < \gamma_0 - r$. Then there exist γ_2, γ' such that $0 < \gamma_2 \leq \gamma_2 + r < \gamma' < \gamma \leq \gamma + r < \gamma_0$ and $s + 1 < \gamma'$. It is easily checked that we have the following properties and their consequences :

- (1) $\mathcal{V}_1(\gamma, \gamma_0)$, therefore $\phi(\cdot)$ has a derivative at $t = 0$ as a \mathcal{B}'_{γ} -valued function (Prop. 8.1),
- (2) $\mathcal{U}_1(\gamma_2, \gamma')$, therefore $P(\cdot)$ has the derivative L_1 at $t = 0$ as a $\mathcal{L}(\mathcal{B}_{\gamma_2}, \mathcal{B}_{\gamma'})$ -valued function (Prop. 6.3),
- (3) $\mathcal{U}_1(\gamma', \gamma_0)$, therefore $P(\cdot)$ has the derivative L_1 at $t = 0$ as a $\mathcal{L}(\mathcal{B}_{\gamma'}, \mathcal{B}_{\gamma_0})$ -valued function (Prop. 6.3),
- (4) $\mathcal{U}_0(\gamma', \gamma)$, therefore $P(\cdot)$ is continuous at $t = 0$ as a $\mathcal{L}(\mathcal{B}_{\gamma'}, \mathcal{B}_{\gamma})$ -valued function (Prop. 6.2).

Lemma 9.4 asserts that $\tilde{v}(\cdot)$ has a continuous derivative, with $\tilde{v}'(0) = im = 0$. The property (2) above ensures that $u(\cdot)$ has a derivative at $t = 0$ as a $\mathcal{B}_{\gamma'}$ -valued function, and that $u'(0)(x) = L_1 1(x) - \tilde{v}'(0) = i \int_G \xi(g, x) d\pi(g) = i\theta(x)$, thus $u'(0) = i\theta$. It follows that $\theta \in \mathcal{B}_{\gamma'}$. Since $\langle \nu, \theta \rangle = im = 0$, Theorem 5.5 shows that there exists a unique $w \in \mathcal{B}_{\gamma'}$ such

that $\langle \nu, w \rangle = 0$ and $(1 - P)w = \theta$, and that w is the sum in $\mathcal{B}_{\gamma'}$ of the series $\sum_{n \geq 0} P^n \theta$. As θ is real valued, so is w . At last, since $\gamma' < \gamma_0 - r$, we have $w \in \mathcal{B}_{\gamma_0 - r}$.

On the basis of the formula of Lemma 9.5, we get

$$\sigma^2 = (\pi \otimes \nu)(\xi^2) - 2i\langle \phi'(0), \theta \rangle = (\pi \otimes \nu)(\xi^2) - 2i\langle \phi'(0), (1 - P)w \rangle.$$

The following lemma allows us to conclude.

Lemma 9.9. *We have*

$$\langle \phi'(0), (1 - P)w \rangle = i(\pi \otimes \nu)(\xi w \circ j).$$

Proof. It is known that, for small $|t|$, $(\lambda(t) - P(t))^* \phi(t) = \phi(t)(\lambda(t) - P(t)) = 0$. Hence, setting $S(t) = \lambda(t) - P(t)$, we have

$$\left(\frac{\phi(t) - \phi(0)}{t} \right) S(t)w + \phi(0) \left(\frac{S(t)w - S(0)w}{t} \right) = \frac{\phi(t)S(t)w - \phi(0)S(0)w}{t} = 0.$$

Observe that $\lambda(t)$ has a derivative at $t = 0$ because the conditions of point (a) in Proposition 9.1 hold. Therefore, since $w \in \mathcal{B}_{\gamma'}$ and $\phi(0) = \nu \in \mathcal{B}'_{\gamma_0}$, the above properties (1) and (4), and then (3) enable us to pass to the limit in the equality. We get

$$\phi'(0)(S(0)w) + \nu(S'(0)w) = 0,$$

or else $\phi'(0)(1 - P)w = \nu[(L_1 - \lambda'(0))w] = \nu(L_1 w) = i \int_M \int_G \xi(g, x) w(gx) d\pi(g) d\nu(x)$. \square

Proof of assertion (b). We know that $\lambda(t) = 1 - \sigma^2 \frac{t^2}{2} + o(t^2)$. Otherwise, since $s + 1 < \gamma_0 - 2r$, there exists η such that $s + 1 < \eta \leq \eta + 2r < \gamma_0$. Since $\mathcal{J}^{\gamma_0 - r}(R, R + S) < +\infty$ and $\mathcal{J}^n(R^2, (R + S)R) < +\infty$ (because $\eta < \gamma_0 - 2r$), the condition $\mathcal{V}_2(\eta, \gamma_0)$ holds. Consequently Proposition 8.2 applies, it follows that $v(\cdot)$, $\phi(\cdot)$, $N(\cdot)$ have second order Taylor's expansions at $t = 0$ as functions with values in $(\mathcal{B}_\eta, N_{\infty, \gamma_0})$, in \mathcal{B}'_η , and in $\mathcal{L}(\mathcal{B}_\eta, \mathcal{B}_{\gamma_0})$ respectively. We get, for all $n \geq 1$,

$$\langle \phi(t), 1 \rangle \langle \mu, v(t) \rangle = 1 + tB + \frac{t^2}{2}C + o(t^2) \quad (A, B \in \mathbb{C})$$

$$\langle \mu, N(t)^n 1 \rangle = \langle \mu, N(0)^n 1 \rangle + t \langle \mu, N_{1,n} 1 \rangle + \frac{t^2}{2} \langle \mu, N_{2,n} 1 \rangle + o_n(t^2).$$

Since $N(0)1 = 0$ and $\varphi_n(t) = \langle \mu, P(t)^n 1 \rangle = \lambda(t)^n \langle \phi(t), 1 \rangle \langle \mu, v(t) \rangle + \langle \mu, N(t)^n 1 \rangle$, with $\lambda(t)^n = 1 - n\sigma^2 \frac{t^2}{2} + o_n(t^2)$, the coefficient b_n of $\frac{t^2}{2}$ in the Taylor expansion of φ_n is $C - n\sigma^2 + \langle \mu, N_{2,n} 1 \rangle$. This enables us to conclude because $\sup_{n \geq 1} \|N_{2,n}\|_{\eta, \gamma_0} < +\infty$. \square

End of the proof of Theorem S'.

Proof of (ii). Let us prove that $\mathbb{E}[(S_n^Z)^2] < +\infty$. Actually, since $2r \leq \gamma_0$, we have, for $k \geq 1$,

$$\mathbb{E}[\xi(Y_k, Z_{k-1})^2] \leq \mathbb{E}[R(Y_k)^2] \mathbb{E}[\psi(Z_{k-1})] = \int_G R^2 d\pi \int_M P^{k-1} \psi d\mu,$$

with $\psi(x) = (1 + d(x, x_0))^{\gamma_0}$. Since $\psi \in \mathcal{B}_{\gamma_0 - 1} \subset \mathcal{B}_{\gamma_0}$, $P \in \mathcal{L}(\mathcal{B}_{\gamma_0})$, and $\mu \in \mathcal{B}'_{\gamma_0}$, we get $\mathbb{E}[\xi(Y_k, Z_{k-1})^2] < +\infty$; hence the claimed property. The function $\varphi_n(\cdot)$ has therefore a

second order derivative at $t = 0$ and $\varphi_n''(0) = -\mathbb{E}[(S_n^Z)^2]$. With the help of Proposition 9.8(b), we obtain $\mathbb{E}[(S_n^Z)^2] = -b_n$, hence $\sigma^2 = \lim_n \frac{1}{n} \mathbb{E}[(S_n^Z)^2]$.

Proof of (i). The method of the proof of Theorem IV.7 of Hennion and Hervé (2001) applies here to the transition probability Q introduced in Section 4.2, yet we give below an adaptation of this method only using P . Set

$$\tilde{\xi} = \xi + w \circ j,$$

where w is the function in Proposition 9.8 (it can be checked that $\tilde{\xi} - Q\tilde{\xi} = \xi$). Recall that $\sigma^2 = (\pi \otimes \nu)[\xi(\xi + 2w \circ j)]$ (Prop. 9.8). From the equality $\xi^2 + 2\xi w \circ j = (\xi + w \circ j)^2 - (w \circ j)^2 = \tilde{\xi}^2 - (w \circ j)^2$, we get

$$\sigma^2 = (\pi \otimes \nu) \left(\tilde{\xi}^2 - (w \circ j)^2 \right).$$

Assume that $\nu(w^2) < +\infty$. Then, using the invariance of ν , we can write

$$\sigma^2 = (\pi \otimes \nu)(\tilde{\xi}^2) - \nu(w^2) = \int_M d\nu(x) \int_G (\tilde{\xi}(g, x)^2 - w(x)^2) d\pi(g).$$

But

$$\int_G \tilde{\xi}(g, x) d\pi(g) = \int_G \xi(g, x) d\pi(g) + Pw(x) = \theta(x) + (w(x) - \theta(x)) = w(x),$$

so that

$$\sigma^2 = \int_M d\nu(x) \int_G (\tilde{\xi}(g, x) - w(x))^2 d\pi(g) = \int_M d\nu(x) \int_G (\xi(g, x) + w(gx) - w(x))^2 d\pi(g).$$

If $\sigma^2 = 0$, we therefore get $\xi(g, x) = w(x) - w(gx)$ $\pi \otimes \nu$ a.e.

To complete the proof of Theorem S', it now suffices to show that the hypothesis $\sigma^2 = 0$ implies $\nu(w^2) < +\infty$. We know that, for all $x \in M$, $w(x) = \sum_{n \geq 0} P^n \theta(x)$. Since $P^n \theta(x) = \mathbb{E}[\theta(R_n x)] = \mathbb{E}[\int \xi(g, R_n x) d\pi(g)] = \mathbb{E}[\xi(Y_{n+1}, R_n x)]$, we have, for all $x \in M$, $w(x) = \lim_n \mathbb{E}[S_n^x]$.

Assume that Z has the distribution ν and that $\sigma^2 = 0$. Then the point (b) of Proposition 9.8 and the fact that $b_n = -\mathbb{E}_\mu[(S_n^Z)^2]$ show that $\sup_n \mathbb{E}[(S_n^Z)^2] = \vartheta < +\infty$. From the inequalities $\int \mathbb{E}[S_n^x]^2 d\nu(x) \leq \int \mathbb{E}[(S_n^x)^2] d\nu(x) = \mathbb{E}[(S_n^Z)^2]$ and Fatou's Lemma, we deduce that $\nu(w^2) \leq \vartheta$. \square

Example : study of σ^2 for sequences of type $(u(Y_n)\chi(Z_{n-1}))_n$.

Suppose that the function ξ is of the form $\xi(g, x) = u(g)\chi(x)$, where u is a non-zero real valued measurable function on G and χ is a real valued locally Lipschitz function on M satisfying $|\chi(x) - \chi(y)| \leq Cd(x, y)(1 + d(x, x_0) + d(y, x_0))^s$. Observe that Condition (RS) holds with $r = s + 1$ and $R(s) = S(g) = C|u(g)|$.

In this context, the next statement based on both Theorem S' and Theorem 5.5 gives a simple sufficient condition for $\sigma^2 > 0$.

Proposition S'. Suppose that the conditions of theorem S hold (with $r = s + 1$ and $R(s) = S(g) = |u(g)|$), that $\int_G u(g)d\pi(g) = 0$, and that $\chi(x) \neq 0$ for some x in the support Σ_ν of the P -invariant measure ν . Then $\sigma^2 > 0$.

Proof. Observe that $m = \pi \otimes \nu(\xi) = 0$. By Theorem S', we shall get $\sigma^2 > 0$ if we prove that there is no real valued function $\tilde{\chi}_1$ in \mathcal{B}_{γ_0-r} such that, for all $x \in \Sigma_\nu$, we have $\xi(g, x) = u(g)\chi(x) = \tilde{\chi}_1(x) - \tilde{\chi}_1(gx) \pi - a.e.$

Let ξ_1 be such a function. Then, by integrating the above equality with respect to the measure π , we get $\tilde{\xi}_1(x) = \int_G \tilde{\xi}_1(gx)d\pi(g) = (P\tilde{\xi}_1)(x)$ for all $x \in \Sigma_\nu$. Since Σ_ν is an absorbing set (for all $x \in \Sigma_\nu$ we have $P(x, \Sigma_\nu) = 1$), this can be rewritten as $\tilde{\xi}_1|_{\Sigma_\nu} = P_{\Sigma_\nu}(\tilde{\xi}_1|_{\Sigma_\nu})$ where P_{Σ_ν} denotes the kernel induced by P on Σ_ν . From $\text{Ker}(P - 1) = \mathbf{C} \cdot 1$ (Th. 5.5), it can be easily proved that the fonctions of \mathcal{B}_{γ_0-r} whose restriction on Σ_ν is P_{Σ_ν} -invariant are constant on Σ_ν . It follows that $\tilde{\xi}_1|_{\Sigma_\nu}$ is constant, thus, for all $x \in \Sigma_\nu$, $u(g)\chi(x) = 0 \pi$ -a.e. This is impossible. \square

REFERENCES

- Bapat R.B. and Raghavan T.E.S. (1997). Non-negative matrices and applications. *Encyclopedia of Mathematics and its applications*, **64**.
- Bhattacharya R. and Majumdar M. (2002). *Economic Theory*, to appear.
- Benda M. (1998). A central limit theorem for contractive stochastic dynamical systems. *J. App. Prob.* **35**, 200-205.
- Bolthausen E. (1982). The Berry-Esseen theorem for strongly mixing Harris recurrent Markov chains. *Z. Wahrsch. verw. Gebiete.* **60**, 283-289.
- Cuny C. (2003). Fonctions harmoniques bornées sur les groupes de Lie. Marches aléatoires et problèmes connexes. *Ann. Inst. H. Poincaré*, to appear.
- Diaconis P. and Freedman D. (1999). Iterated random functions. *SIAM Rev.* **41**, no. 1, 45-76.
- Dudley R.M. (1989). Real analysis and probability. *Wadsworth, Pacific Grove, CA*.
- Duflo M. (1997). Random Iterative Models. *Applications of Mathematics, Springer-Verlag Berlin Heidelberg*.
- Dunford N. and Schwartz J.-T. (1958). Linear operator, Part I. *Pure and Applied Mathematics. Vol. VII. Interscience*.
- Hennion H. (1991). Dérivabilité du plus grand exposant caractéristique des produits de matrices aléatoires indépendantes à coefficients positifs. *Ann. Inst. Henri Poincaré, Vol. 27, No 1, p. 27-59*.

- Hennion H. (1993). Sur un théorème spectral et son application aux noyaux lipchitziens. *Proceeding of the A.M.S*, vol. 118, No 2, pp. 627-634.
- Hennion H. (1997). Limit theorems for products of positive random matrices. *The Annals of Probability*, Vol. 25, No 4, p. 1545-1587.
- Hennion H. and Hervé L. (2001). Limit theorems for Markov chains and stochastic properties of dynamical systems by quasi-compactness. *Lecture Notes in Mathematics No 1766*, Springer.
- Keller G. and Liverani C. (1999). Stability of the Spectrum for Transfer Operators. *Ann. Scuola Norm. Sup. Pisa. Cl. Sci. (4) Vol. XXVIII*, pp. 141-152.
- Le Page E. (1983). Théorèmes de renouvellement pour les produits de matrices aléatoires. Equations aux différences aléatoires. *Séminaires de Rennes*.
- Le Page E. (1989). Régularité du plus grand exposant caractéristique des produits de matrices aléatoires indépendantes et applications. *Ann. Inst. Henri Poincaré*, Vol. 25, No 2, p. 109-142.
- Meyn S.P. and Tweedie R.L. (1993). Markov chains and stochastic stability. *Springer-Verlag, Berlin*.
- Milhaud X. and Raugi A. (1989). Etude de l'estimateur du maximum de vraisemblance dans le cas d'un processus auto-régressif : convergence, normalité asymptotique, vitesse de convergence. *Ann. Inst. H. Poincaré*, Vol. 25, No 4, pp. 383-428.
- Nagaev S.V. (1957). Some limit theorems for stationary Markov chains. *Theory Probab. Appl.*, **11**, no. 4, 378-406.
- Peigné M. (1993). Iterated function schemes and spectral decomposition of the associated Markov operator. *Séminaires de Probabilité de Rennes*.
- Pollicott M. (2001). Contraction in mean and transfer operators. *Dyn. Syst*, Vol. 16, No 1, pp. 97-106.
- Wei-Biao Wu and Woodroffe M. (2000). A central limit theorem for iterated random functions. *J. App. Prob.* **37**, 748-755.